



Dépôt institutionnel  
Repository

Institut de recherche  
Robert-Sauvé en santé  
et en sécurité du travail

Ceci est la version pré-publication, révisée par les pairs, de l'article suivant :

Bakhiyi, B., Gravel, S., Ceballos, D., Flynn, M. A. et Zayed, J. (2018). Has the question of e-waste opened a Pandora's box? An overview of unpredictable issues and challenges. *Environment International*, 110, 173-192.

La version finale de l'article est disponible à <https://doi.org/10.1016/j.envint.2017.10.021>.

Cet article peut être utilisé à des fins non commerciales.

**Avis :** L'IRSST encourage son personnel scientifique et tout chercheur dont il finance en tout ou en partie les travaux ou qui bénéficie de son programme de bourses à faire en sorte que les articles issus de ces travaux soient librement accessibles au plus tard un an après leur publication dans une revue savante.

<https://www.irsst.qc.ca/Portals/0/upload/5-institut/politiques/Libre-acces.pdf>

[communications@irsst.qc.ca](mailto:communications@irsst.qc.ca)

---

This is the accepted manuscript peer reviewed version of the following article:

Bakhiyi, B., Gravel, S., Ceballos, D., Flynn, M. A., & Zayed, J. (2018). Has the question of e-waste opened a Pandora's box? An overview of unpredictable issues and challenges. *Environment International*, 110, 173-192.

It is available in its final form at <https://doi.org/10.1016/j.envint.2017.10.021>.

This article may be used for non-commercial purposes.

**Disclaimer:** The Institut de recherche Robert-Sauvé en santé et en sécurité du travail (IRSST) encourages its scientific staff and all researchers whose work it funds either in whole or in part or who benefit from its scholarship program to ensure that the articles resulting from their work be made publicly accessible within one year of their publication in a scholarly journal.

<https://www.irsst.qc.ca/Portals/0/upload/5-institut/politiques/Open-access.pdf>

[communications@irsst.qc.ca](mailto:communications@irsst.qc.ca)

**Has the question of e-waste opened a Pandora's box? An overview of unpredictable issues and challenges**

Bouchra Bakhiyi<sup>a</sup>, Sabrina Gravel<sup>a, b</sup>, Diana Ceballos<sup>c</sup>, Michael A. Flynn<sup>d</sup>, Joseph Zayed<sup>a, b</sup>

<sup>a</sup> Department of Environmental and Occupational Health, School of Public Health, Université de Montréal, Montreal, Quebec, Canada

<sup>b</sup> Institut de recherche Robert-Sauvé en santé et en sécurité du travail (IRSST), Montreal, Quebec, Canada

<sup>c</sup> Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA, USA

<sup>d</sup> National Institute for Occupational Safety and Health, Cincinnati, OH, USA

**Keywords:**

E-waste management, chemical contaminants, electronic recycling, legacy contamination, human health, occupational health

**Corresponding author:**

Joseph Zayed, Ph.D.

Adjunct Professor

Department of Environmental and Occupational Health

School of Public Health

University of Montreal

P.O. Box 6128, Main Station

Montreal, Quebec

Canada, H3C 3J7

Telephone: +1 514 343 5912

Fax: +1 514 343 2200

Email: joseph.zayed@umontreal.ca

## Abstract

Despite regulatory efforts and position papers, electrical and electronic waste (e-waste) remains ill-managed as evidenced by the extremely low rates of proper e-waste recycling (e-recycling) worldwide, ongoing illegal shipments to developing countries and constantly reported human health issues and environmental pollution. This review aims to expose the complexity of e-waste problems and suggest possible upstream and downstream solutions. Exploring e-waste issues is akin to opening a Pandora's Box. Thus, a review of prevailing e-waste management practices reveals complex and often intertwined gaps, issues and challenges. These include the absence of any consistent definition of e-waste to date, a prevalent toxic potential still involving already banned or restricted hazardous components such as heavy metals and persistent and bioaccumulative organic compounds, a relentless growth in e-waste volume fueled by planned obsolescence and unsustainable consumption, problematic e-recycling processes, a fragile formal e-recycling sector, sustained and more harmful informal e-recycling practices, and more convoluted and unpredictable patterns of illegal e-waste trade. A close examination of the e-waste legacy contamination exposes critical human health concerns, including significant occupational exposure during both formal and informal e-recycling, and persistent environmental contamination, particularly in some developing countries. However, novel e-waste contaminants as well as unexpected sources and environmental fates of contaminants are among the emerging issues that raise concerns. Moreover, scientific knowledge gaps remain regarding the complexity and magnitude of the e-waste legacy contamination, specifically, a comprehensive characterization of e-waste contaminants, information on the scale of legacy contamination in developing countries and on the potential environmental damage in developed countries, and a stronger body of evidence of adverse health effects specifically ascribed to e-

waste contaminants. However, the knowledge accumulated to date is sufficient to raise awareness and concern among all stakeholders. Potential solutions to curb e-waste issues should be addressed comprehensively, by focusing on two fronts: at the upstream and downstream levels. Potential upstream solutions should focus on more rational and eco-oriented consumer habits in order to decrease e-waste quantities while fostering ethical and sustained commitments from manufacturers, which include a limited usage of hazardous compounds and an optimal increase in e-waste recyclability. While progressively reducing illegal e-waste trade, downstream solutions should introduce stronger reverse logistics, together with upgraded, more affordable, and eco-friendly and worker friendly e-recycling technologies to ensure that benefits are derived fully and safely from the great economic potential of e-waste.

## Contents

<b>1. Introduction.....</b>	<b>4</b>
<b>2. Background: a bird's-eye view of the history of e-waste management.....</b>	<b>5</b>
<b>3. Ongoing gaps, issues and challenges in e-waste management.....</b>	<b>9</b>
<b>4. E-waste legacy contamination.....</b>	<b>16</b>
4.1. Legacy contamination and human body burden .....	22
4.2. Remaining gaps in scientific knowledge in legacy contamination .....	25
<b>5. Potential solutions to curb e-waste issues.....</b>	<b>28</b>
<b>6. Conclusion .....</b>	<b>34</b>
<b>Acknowledgments .....</b>	<b>35</b>
<b>References.....</b>	<b>36</b>
<b>APPENDIX.....</b>	<b>59</b>

## 1. Introduction

Electronic waste (e-waste), or waste electrical and electronic equipment (WEEE), among other terms are terms used to cover electrical and electronic equipment discarded as waste without intent of reuse (Step, 2014). E-waste offers particularly high potential for recovery as it contains valuable recyclable components such as gold, platinum and silver. However, e-waste also contains non-negligible amounts of potentially toxic substances (e.g., cadmium and lead) and is thus considered hazardous when improperly managed. These findings highlight the need for the safe and smart management (including proper recycling) and commercial exploitation of e-waste while preserving human health and environmental integrity, given the large volume of e-waste generated worldwide annually (41.8 million metric tonnes (MMT) estimated in 2014) and its fast growth (Lundgren, 2012; Kiddee et al., 2013; Cucchiella et al., 2016; Step, 2016, Baldé et al., 2015). Canada, which generated an estimated 725,000 tonnes of e-waste in 2014, is well below the 2014 top five e-waste global generators, which were the United States (7.1 MMT), China (6.0 MMT), Japan (2.2 MMT), Germany (1.8 MMT) and India (1.7 MMT). However, with 20.4 kg of e-waste generated annually per inhabitant, Canada remains one of the highest contributors to e-waste volume in relative quantities in the Americas, right between the United States (22.1 kg/inh.) and the Bahamas (19.1 kg/inh.) (Baldé et al., 2015).

This review aims to expose the complexity of e-waste problems. Its objectives are (a) to provide an brief overview of the historical aspects of e-waste management; (b) to identify gaps, issues and challenges that greatly complicate e-waste management; (c) to gain insight into the current e-waste legacy contamination in terms of critical, emerging or still-unknown human health issues (including occupational health

concerns) and environmental contamination and (d) to propose solutions that could potentially curb e-waste issues both upstream and downstream.

## **2. Background: a bird's-eye view of the history of e-waste management**

In the 1970s and 1980s, hazardous waste, including e-waste, was commonly shipped from industrialized countries to less developed nations in Asia, Africa, Central America and Eastern Europe (UNEP, 2010). The hazardous waste trade is rooted in the “Not in My Back Yard” syndrome in developed countries, an expression of the public’s vehement stand against poor management of hazardous waste, including e-waste. Since the 1970s, it has led to the adoption of more stringent laws in the developed countries, such as The Resource Conservation and Recovery Act (RCRA) in the United States in 1976 (UNEP, 2010), which resulted in escalating the costs of hazardous waste disposal, while these costs remained low in less developed countries (Massari and Monzini, 2006; Andrews, 2009; UNEP, 2010).

E-waste trading led to heavy environmental contamination in receiving countries, where primitive recycling methods, incineration and landfilling of hazardous waste were widely practiced, supported by inadequate environmental awareness, controls and regulations (UNEP, 2010). To fight what was called the “toxic trade,” the Basel Convention on the Control of the Transboundary Movements of Hazardous Wastes and their Disposal was adopted in 1989 and came into force in 1992. The aim of this international treaty is to regulate the export of hazardous waste from industrialized countries (called “Annex VII countries” and composed of parties to the Basel Convention that are members of the Organization of the Economic Co-operation and Development (OECD) or the European Union (EU), as well as Liechtenstein) to less developed and vulnerable nations (called “non-Annex VII countries” and composed of all other

parties to the Basel Convention). The fundamental purpose of the Basel Convention is to promote safe and sound hazardous waste management in order to safeguard human health and the environment. Its main objectives also include the limitation of hazardous waste generation and the restriction of hazardous waste exports unless the receiving country has confirmed the existence of environmentally sound practices for managing the imported waste (Andrews, 2009; Ahmad Khan, 2016).

Amendments to the Basel Convention, known as the Basel Ban or the Ban, were adopted in 1995 and 1997 to completely prohibit the export of hazardous wastes from Annex VII countries to non-Annex VII countries, while Annex VIII was added in 1998 to include e-waste. However, Annex IX, also added in 1998, still allows the export of certain categories of e-waste for strict reuse, for the purpose of giving the receiving parties access to the digital world through second-hand equipment (Basel Convention, 2011; Ahmad Khan, 2016).

A series of regulations and policies have since emerged worldwide at the regional, national and global levels to promote reuse and proper recycling, as well as a reduction in the use of toxic raw materials. For example, in 2003, the European Union adopted significant regulations such as the WEEE Directive, which sought to enhance e-waste collection, reuse and recycling, and the Restriction of the Use of Certain Hazardous Substances in Electrical and Electronic Equipment (RoHS) Directive, requiring substitutions for or the limitation of certain toxic substances, including heavy metals and flame retardants (European Commission, 2017).

On a global scale, the Nairobi Declaration on the Environmentally Sound Management of Electrical and Electronic Waste was launched in 2006 (Lundgren, 2012). A multi-stakeholder partnership known as the

Solving the E-waste Problem (StEP) Initiative, involving United Nations agencies as well as academic and governmental organizations, among others, was instituted in 2007 to achieve more sustainable e-waste management through an international information-sharing platform (Sthiannopkao and Wong, 2013). An entire battery of tools has also been developed and adopted over time in developed and some developing countries to support the safe and optimal handling of the e-waste stream: Extended Producer Responsibility (EPR), Life Cycle Assessment (LCA), Material Flow Analysis (MFA), Multi-Criteria Analysis (MCA) (Kiddee et al., 2013).

Unfortunately, despite regulatory efforts and position papers, e-waste is still ill-managed, with the proper e-waste recycling (e-recycling) rate remaining extremely low worldwide, at roughly 15.5% of the global volume generated in 2014 (Baldé et al., 2015). Even more alarming is the fact that illegal shipments to more vulnerable countries continue to abound, as disclosed by a report of the Interpol Pollution Crime Working Group (Interpol, 2009).

Through a 2014-2016 investigation, a US environmental watchdog group called the Basel Action Network (BAN) unveiled the continuing traffic of e-waste from the US to developing countries (mostly China) and involving computer manufacturers, certified recycling companies and at least one major charity organization. More than 90% of e-waste was actually exported illegally under the guise of second-hand equipment (Hopson and Puckett, 2016). Following the BAN report, the US Environmental Protection Agency (US EPA) strongly condemned such illegal activities and levied a severe fine on of the US electronic recyclers involved (US EPA, 2016a; WDE, 2016).



Given the increasing volume of e-waste being generated worldwide, it can be assumed that illegal shipments to the developing world have been rising steadily since the Basel Convention, leading to a globalization of e-waste issues. About 50-80% of the e-waste generated in developed countries is considered to be illegally exported to low- and middle-income countries (Ghosh et al., 2016; Someya et al., 2016). China still receives the lion's share of all illegal e-waste, although countries such as the Philippines, India, Nigeria and Ghana remain attractive destinations. It is worth mentioning that most of the receiving countries have nonetheless ratified the Basel Convention and the Basel Ban (Rucevska et al., 2015; Hopson and Puckett, 2016; Terazono et al., 2017). It is estimated that between 1.5 and 2 MMT of e-waste are exported illegally to China from the European Union each year, despite the import ban imposed by China on all e-waste in 2000 (Huisman et al., 2015; Geeraerts et al., 2016). Moreover, a recent survey conducted in Hong Kong revealed that some traders do not hesitate to "re-export" 90-100% of the used televisions shipped from Japan to China (Sugimura and Murakami, 2016).

Human health risks and environmental damage, particularly when related to unsafe e-waste handling and processing in developing countries, have therefore raised substantial concerns in the scientific community. Through a systematic review, some authors have found what they consider to be plausible detrimental effects associated with exposure to e-waste contaminants. The vulnerable populations identified include not only workers, but also children and pregnant women living within or in the vicinity of unregulated recycling sites. E-waste has therefore become an emerging health risk for critically exposed populations in developing countries (Chen et al., 2011; Grant et al., 2013). This finding is attributed in particular to the relentless growth in e-waste and the ensuing and constantly reported environmental contamination with persistent and bioaccumulative organic pollutants, including flame retardants (Song and Li, 2015;

Awasthi et al., 2016). Occupational health and safety issues related to both the proper recycling sector and unregulated e-waste activities are now subjected to greater scientific scrutiny. However, many aspects of the potential occupational exposure remain unknown (Ceballos and Dong, 2016; Tue et al., 2016a).

### **3. Ongoing gaps, issues and challenges in e-waste management**

For e-waste management to be respectful of the environment and of human health, it has to be properly and effectively handled. Effective e-waste management first consists of collecting and sorting e-waste, repairing and reusing it whenever possible. Then, end-of-life e-waste is processed to remove and decontaminate all potentially toxic compounds, to properly recover valuable materials, and finally, to safely dispose of toxic parts and non-recyclable residuals (Namias, 2013; Baldé et al, 2015; Step, 2016). Unfortunately, e-waste management is currently facing significant, varied and complex gaps, issues and challenges that are often intertwined and that make e-waste problems so arduous to resolve.

One of the most critical gaps is the absence of a consistent definition of e-waste, which makes it even harder to precisely assess the massive volume generated worldwide and the dynamics of the current illegal e-waste trade, which in turn rank among the main e-waste issues (Lepawsky, 2015; Ahmad Khan, 2016).

The evolving nature of e-waste, attributed among others to technological progress, challenges the creation of a harmonized and definitive list of end-of-life electrical and electronic products (Ahmad Khan, 2016; Magalini, 2016). Hence, the WEEE Directive was amended in 2012 to include photovoltaic panels (European Commission, 2017).

The relentless manufacture of new electrical and electronic products (referred to as e-products in this paper), and the continuous growth of e-waste volume is also in itself a considerable issue. With an

estimated annual growth rate of 4-5%, a global production of close to 50 MMT is predicted for 2018 (Baldé et al., 2015), which is plausible given that more than 56 MMT of e-products came on the market in 2012 (Honda et al., 2016). For China alone, Zeng Xianlai et al. (2016) suggested an annual growth of 25.7%, which could generate 15.5 MMT of e-waste in 2020, and almost double in 2030. This continuous growth is mainly fueled by planned obsolescence which refers to the continuing decrease in the lifetime of e-products as a result of rapid technological updates, coupled with a drastic reduction in retail prices, and strongly backed by marketing and advertising. E-products are quickly perceived as outdated, and undergo premature equipment failure (Pickren, 2015; Echegaray, 2016; LeBel, 2016; Zeng et al., 2017). This is particularly so for small electronic devices such as personal computers (PC), whose lifespan shortened from 4-6 years in 1997 to two years in 2005 (Lundgren, 2012), and cellular phones, whose lifespan collapsed in the United States from approximately 30 months in 1995 to less than 20 months in 2005 (Ryneal, 2016).

More often than not, defective or damaged e-products tend to be replaced rather than repaired, further weakening e-products recovery and repair sector. This is in part due to limited public interest, technological obsolescence leading to the unavailability of spare parts, unsuitable tools, predominance of self-learning among repairers, time-consuming tasks during the repair process, high labor costs, and lack of strong initiative and effective logistics for testing and repairing defective items (Echegaray, 2016; Sabbaghi et al., 2017). The design itself of e-products discourages their repair as their components are hastily assembled and securely attached (e.g., glued, bolted, or soldered) and thus repair becomes extremely difficult and time-consuming, if possible at all (Lundgren, 2012; Pickren, 2015; Sabbaghi et al., 2017). Obsolescence of e-products can however also be subjective or psychological, leading to

unsustainable consumption of e-products. Echegaray (2016) defined subjective obsolescence as a consequence of “*the subjective devaluation of product perception based on learned experience, emotional attachments or benefits, status achievement, fashion, or esthetic quality.*” The author found that consumers adopt throwaway practices with ease, replacing e-products because of new technology appeal, aesthetics or changing fashions, rather than declining performances or technical failure. This frenetic consumerism may also be stimulated by a growth in buying power, as shown by Kumar et al. (2017) who identified a positive correlation between the Gross Domestic Product (GDP) and e-waste generation in any country in the world, regardless of the number of inhabitants.

Not only do we witness the growth of e-waste volume, but the process of calculating global e-waste volume also faces challenges due to the aforementioned absence of a clear e-waste definition, the extreme scarcity and inefficient collection of official data, the scanty data on both legal and illegal e-waste trans-boundaries and the lack of awareness among stakeholders (Honda et al., 2016; Morris and Metternicht, 2016; Petridis et al., 2016; Tran et al., 2016; Kumar et al., 2017). These data remain fundamental for receiving countries, as e-waste volume is generated not only by locally purchased e-products but also by imported second-hand equipment (referred to as the “*invisible inflow*”).

The dearth of reliable data, combined with an increased e-waste volume, give rise to unpredictable and convoluted patterns of illegal e-waste trade. Geographies and the dynamics of e-waste trafficking have turned it into a more complex web. For instance, despite the official ban on e-waste imports, several Asian countries exchange e-waste between themselves, and some countries such as Ghana and Nigeria are still the main receivers of exports not only from Israel, Belgium, the United States and Canada but also from Singapore and India (Lepawsky, 2015; Lines et al., 2016; Someya et al., 2016). This could be in part

228 attributed to some gaps, limitations and contradictions that can be highlighted in the Basel Convention.  
229 Notably, the coming into force of the Basel Ban is still delayed, as only 89 countries have ratified it so far  
230 (Basel Convention, 2017) while 90 ratifications are required (Ahmad Khan, 2016). Major e-waste  
231 producing countries like the United States have not yet ratified the Basel Convention itself, although the  
232 US was among the first signatories (Bradford, 2011; Ahmad Khan, 2016). In addition, by allowing e-  
233 waste exportation for strict reuse, Annex IX has provided fertile ground for the illegal trade through re-  
234 categorization of non-functional or nearly-out-of-date e-waste. False custom declarations are known to be  
235 common, disguising hazardous e-waste as used e-products for the second-hand market, plastic or metal  
236 scrap (Ahmad Khan, 2016; Geeraerts et al., 2016; Milovantseva and Fitzpatrick, 2016). There are also  
237 certain gaps in the customs rules of several countries, including a shortage of international information  
238 sharing, human resources, training and logistics to check each container received and to distinguish  
239 between still-functional goods and actual waste. Moreover, hazardous waste, non-hazardous waste and  
240 non-waste (used and new goods) often have identical six-digit-codes in the Harmonized Commodity  
241 Description and Coding System (HS) used by customs services across the globe (Ahmad Khan, 2016,  
242 Huisman et al., 2015; Geeraerts et al., 2016; Grant and Oteng-Ababio, 2016).

243 Another pivotal issue in e-waste management is the inherent difficulties of the e-recycling processes.  
244 Despite some significant improvements, metal recovery technologies are still underperforming (e.g.,  
245 difficulties in recycling single valuable metals), immature and/or polluting (Zhang and Xu, 2016). Some  
246 e-waste components such as printed circuit boards, CRT monitors and LCD screens also remain extremely  
247 hard to recycle mainly due to their complexity, their hazardous nature and inefficient recycling  
248 technologies, so much so that many recyclers simply refuse to process them (Edwards, 2016; Singh et al,

2016). The same can be noticed with e-products of a more compact design, for which recycling call for a state-of-the-art, costly, and labour- and energy-intensive processes to safely segregate and optimally recover materials. However, few facilities are equipped with the technical know-how required to end-process e-waste (Lundgren, 2012; Duan et al., 2016; D'Adamo et al., 2016; Zhang and Xu, 2016; Tansel, 2017).

Perhaps the most important problem in terms of e-waste's impact on the environment and human health is the fact that numerous e-waste products still contain hazardous components that have already been banned or restricted in several countries. These include heavy metals, ozone-depleting substances and persistent organic pollutants (POPs) such as polychlorinated biphenyls (PCBs) and brominated flame retardants (Baldé et al., 2015; Zheng Xiaobo et al., 2016; Matsukami et al, 2017). Thus, the global e-waste generated in 2014 was estimated to contain 2.2 MMT of leaded glass, 0.3 MMT of lead/lithium/cadmium/mercury-based batteries and 4,000 tons of ozone-depleting substances such as chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs) (Baldé et al., 2015).

There have been efforts to substitute potentially toxic substances in the manufacturing process of e-products. For instance, flat screen monitors with liquid crystal displays (LCDs) have replaced leaded-glass CRT monitors. However, their recycling is also hazardous due to the presence of toxic chemicals such as Hg, As, Cr and Ba (US EPA, 2016b; Woo et al., 2016; Tansel, 2017). A similar issue is observed with the substitution of certain chemicals in the manufacturing process, like the replacement of PBDEs by halogen-free organophosphorus flame retardants, where the substituent is alas proving to be more persistent than predicted (Gramatica et al., 2016). The promising development of green electronics processes, such as the inclusion of biodegradable electronic components (e.g., cellulose-based printed circuit boards), which

would reduce the toxic potential of e-waste, is still unfortunately fairly slow (Guna et al., 2016; Ma et al., 2016). Additionally, there are limitations in the provided information regarding the chemicals included in e-products, making the proper disposal of toxic waste even more challenging. Indeed, the publicly available and/or relevant data for recyclers is scarce, and there is a deficient chemical composition data disclosure between stakeholders involved in the e-product chain (e.g., chemical and formulation manufacturers, brand-name owners, recyclers) (Scruggs et al., 2016; US EPA, 2016b; Woo et al., 2016). Finally, the e-waste recycling industry often employs individuals from socially marginalized or vulnerable groups such as temporary workers, immigrants or prisoners (Ceballos and Dong, 2016). In developing countries, it is not uncommon to find children working in e-recycling (Robinson, 2009). For example, in Guiyu, China e-waste workers tend to be internal migrants, many of whom are minors (Sthiannopkao and Wong, 2013). An evaluation of an electronic scrap recycling facility in the United States found that roughly half of all employees were among immigrants (Page et al., 2015). Similarly, the United States Department of Justice (2010) investigated safety concerns of an electronic waste recycling program among inmates in Federal Prisons in the United States. Workers from marginalized social groups frequently have less economic security and their vulnerable social position often contributes additional barriers to safety and health at work which leads to increased risk for occupational injuries and illness and aggravates the safety challenges faced by the e-waste industry (Krieger, 2010; Flynn, 2014).

All of these interconnected gaps, issues and challenges mentioned here render the formal e-recycling sector<sup>1</sup> more fragile. While most e-waste is handled in an informal manner in developing countries (Gu et

---

<sup>1</sup> Formal e-waste recycling refers to activities carried out in facilities that are usually licensed and/or under the control of some legal authority and that comply with environmental laws and regulations. (Fujimori et al., 2012; McCann and Wittmann, 2015; Ceballos and Dong, 2016).

al., 2016), formal e-recycling rates of the total e-waste remain very low in developed countries: 35% in the European Union in 2012 (Huisman et al., 2015), 29% in the United States in 2012 (Seeberger et al., 2016) and around 24 to 30% in Japan, 12% in Canada and roughly 1% in Australia for all e-waste generated in 2014 (Baldé et al., 2015). Formal e-recycling is usually seen as unprofitable due to the need for high investment, dependency on governmental subsidies and elevated labour costs, all of which are outweighed by underperforming material recovery and low profits (Lundgren 2012; Ryneal, 2016). There is also a lack of regulation, as well as challenging certification programs, limited implementation of various WEEE directives, and a great need of federal government involvement in compliance measures and financial support (Mohanty et al., 2015; Edwards, 2016; Morris and Metternicht, 2016). In addition, many deficiencies remain in official collection systems in developed countries, in part because of the lack of collection points, the financial costs associated with recycling and transportation, and even stealing activities (e.g., illegal traffic in e-waste) (Huisman et al., 2015; Li et al., 2015; McCann and Wittmann, 2015; Garrath et al., 2017). On top of that, massive e-waste quantities and the fast growth place a particular burden on collection logistics and proper e-recycling infrastructures (Tansel, 2017). In developing countries, informal collection systems are in most cases well-organized, offering more convenient door-to-door services at better prices than offered by formal collectors (Gu et al., 2016; Borthakur and Govind, 2017). Sustainability of informal e-recycling activities in developing countries can be explained by the fact that they provide convenient livelihood system through the business and job opportunities they create,

---

By contrast, informal e-recycling is generally considered less appropriate as it is scantily regulated, is potentially more harmful, has a very limited investment capacity for final disposal and usually takes place in small-scale and scattered workshops or domestic backyards (McCann and Wittmann, 2015; Ceballos and Dong, 2016; Tao et al., 2016; Terazono et al., 2017).



thus improving the quality of life of vulnerable populations (Ahmad Khan. 2016; Ardi and Leisten, 2016; Grant and Oteng-Abadio, 2016). Despite government initiatives, the implementation of a formal e-recycling sector in developing countries faces many barriers, such as high costs, supply deficiency, and the lack of strict regulation, a clear legal framework, adequate infrastructure, specific training, risk awareness and financial incentives (McCann and Wittman, 2015; Geeraerts et al., 2016; Gu et al, 2016). Finally, another crucial issue lay in the inconsistency in public awareness and willingness regarding proper e-recycling practices. Level of awareness varies from a complete lack among consumers in some developing countries (Liang and Sharp, 2016; Nartey, 2016) to favourable attitudes toward, but marginally adequate behaviour regarding formal e-recycling, in both developed and developing countries (Borthakur and Govind, 2017; Echegaray and Hansstein, 2017).

#### **4. E-waste legacy contamination**

E-waste handling and processing have the potential to release a complex blend of contaminants in various environmental matrices. These releases can be more or less significant, depending on the processes used and the protective measures implemented. Contaminants in e-waste are released in highly heterogeneous mixtures, whose composition varies according to e-waste types, age, as well as handling and processing. The physical state of the released contaminants varies with the nature of the handling process, and includes particulate matter, gas, vapour, aerosol, solid residue left after a smelting or leaching process, liquid (spent acid or waste water) or semi-liquids (sludge from leaching solutions). Three large categories of contaminants may be discharged (Schluep et al., 2009; Lundgren, 2012; Cayumil, 2016):

- different classes of primary contaminants, mostly metals and persistent and bioaccumulative organic substances as part of e-waste composition (Table 1), generally during pre-processing activities (e.g., manual dismantling, shredding, and crushing), end-processing activities (e.g., open burning of e-waste or pyrometallurgical/ hydrometallurgical routes for extracting precious materials), and incineration of e-waste (both integral and residual), and as landfill leachates;
- secondary contaminants, mostly persistent and bioaccumulative organic substances (Table 2) that derive from the combustion of primary contaminants such as PCBs or other ingredients (e.g., polyvinyl chloride/PVC in plastic or cabling), or during e-waste open burning, smelting or incinerating operations;
- tertiary contaminants, mostly acids and cyanides (Table 2), emitted by chemicals such as the acids and cyanides used in material recovery processes.

**Table 1**

**Major e-waste primary contaminants**

Contaminants	Example of sources	Main types of toxicity
<b>METALS</b>		
Aluminum	WPCBs, microchips, hard drives, LED monitors, plastic housing; plastics, cables and wires containing inorganic flame retardants (e.g. aluminum hydroxide and trihydroxide)	Lung irritant, neurotoxic
Americium 241	Ionization smoke detectors	Carcinogenic, in the case of high dose radiation
Antimony	Tin-lead alloys, WPCBs, CRT; LCD TVs; plastics, cables and wires containing inorganic flame retardants (e.g. antimony trioxide <sup>1</sup> )	Lung, eye and gastro-intestinal irritant
Arsenic	Dopant for semi-conductors, PTVs, LCD monitors and TVs	Carcinogenic, hematotoxic, endocrine disrupter
Barium	CRT, fluorescent lamps, LCD TVs, PTVs, gutters in vacuum tubes.	Neurotoxic, cardiotoxic, gastro-intestinal irritant
Beryllium	WPCBs, power supply boxes, ceramic components	Berylliosis, carcinogenic

Contaminants	Example of sources	Main types of toxicity
Cadmium	Batteries, toners, cartridges, plastics, WPCBs, solder, chip resistors, CRT, PTVs, cell phones, infrared detectors	Carcinogenic, cardiotoxic, nephrotoxic, endocrine disrupter
Cobalt	Batteries, hard drives, laptop computers, LCD monitors and TVs, PTVs, CRT	Cardiotoxic, allergen (asthma), possibly carcinogenic to human (IARC)
Copper	Cables, electrical wiring, WPCBs, microprocessors, terminal strips, plugs, PTVs, cell phones	Lung, eye and gastro-intestinal irritant
Gallium	Data tapes, integrated circuits	Skin and probably eye and mucous membrane irritant
Hexavalent Chromium VI	Corrosion resistant coatings, WPCBs, data tapes, floppy disks, pigments, PTVs	Carcinogenic (lung cancer), sensitizer, skin irritant
Indium	LCD TVs, transistors, rectifiers	Probably carcinogenic to human (IARC <sup>2</sup> ), eye and lung irritant, indium lung disease
Lead	CRT (glass, solder), LCD TVs, PTVs, fluorescent tubes, WPCBs, lead-acid batteries	Probably carcinogenic to human (IARC <sup>2</sup> ), neurotoxic, cardiotoxic, nephrotoxic, endocrine disrupter
Lithium	Rechargeable batteries	Skin and eye irritant
Manganese	Cell phones, CRT, PCBs	Cardiotoxic, neurotoxic, lung irritant, endocrine disrupter <sup>3</sup>
Mercury	Fluorescent tubes, compact fluorescent lamps, batteries, switches, thermostats, sensors, monitors, LCD TVs, laptop computers	Neurotoxic, skin, eye and gastro-intestinal irritant, endocrine disrupter
Nickel	Nickel-cadmium batteries, ceramic components of electronics, computers, LCD monitors and TVs, laptop computers	Carcinogenic, sensitizer
Platinum	WPCBs, LCD and LED notebooks, cell phones	Eye and lung irritant
Selenium	Rectifiers, WPCBs, old photocopiers	Eye, skin and lung irritant, selenosis at high concentrations
Silver	PTVs, laptop computers, LCD and LED monitors	Nephrotoxic, reprotoxic
Thallium	Semi-conductors, batteries	Neurotoxic, cardiotoxic, hepatotoxic, birth defects
Tin	LCD screens, tin-lead alloys, LED monitors	Skin and eye irritant, hematotoxic, hepatotoxic
Tungsten	LCD and LED monitors	Lung, eye and skin irritant
Vanadium	PCBs, red phosphor emitters, tablet PCs	Cardiotoxic, nephrotoxic, skin and lung irritant
Yttrium	LCDs, superconductors, lasers	Lung and eye irritant
Zinc	WPCBs, PTVs, CRTs, batteries, soldering flux, cell phones, plastics, wire and cables containing inorganic flame retardants (e.g., in the form of zinc stannate)	Neurotoxic, hematotoxic, gastric irritant, may induce fume fever (inhalation of large amount of zinc in fume or dusts), probably endocrine disrupter <sup>4</sup>

## ORGANIC POLLUTANTS

<b>Halogenated flame retardants</b>	WPCBs, plastics	Endocrine disrupter, neurotoxic,
- Brominated flame retardants (e.g., PBDEs, TBBPA, HBCD)	Condensers, transformers, plastics	carcinogenic (chlorinated flame retardants)
- Novel brominated flame retardants (e.g., DBDPE, TBPH, TBBPA-BGE)		

Contaminants	Example of sources	Main types of toxicity
- Chlorinated flame retardants (e.g., DPs, DBHCTD)		
<b>Halogen-free flame retardants</b> Organophosphorus-based flame retardants <sup>5</sup> (e.g., TCEP, TCIPP, TDCIPP, TPHP/TPP, BPA-BDPP, PBDPP, and DOPO) Nitrogen-based flame retardants <sup>6</sup> (e.g., melamine cyanurate, melamine polyphosphate)	IT housing, plastics, epoxy resins in WPCBs	<b>Organophosphorus:</b> Endocrine disrupter  <b>Nitrogen-based:</b> Nephrotoxic, neurotoxic
PCBs	Old capacitors and transformers, fluorescent lamps, electrical motors	Endocrine disrupter, hepatotoxic, carcinogenic
<b>OZONE-DEPLETING SUBSTANCES</b>		
E.g., CFCs, HCFC, HFC, HCs	Old refrigerators, freezers and air conditioning units, insulation foam	Neurotoxic, lung and eye irritants
<b>OTHER COMPONENTS</b>		
Phthalates	Plasticizers to soften plastics and rubber	Endocrine disrupter, reprotoxic, hepatotoxic
PFOS/F	Antireflective coating	May effect lipid metabolism (increase in blood cholesterol level)
PVC	Wiring and computer housing	Related to toxicity of dioxins and furans generated during PVC burning (see Table 2)

<sup>1</sup> Often associated with other halogenated and non-halogenated flame retardants (Aschberger et al., 2017)

<sup>2</sup> According to the International Agency for research on cancer (IARC) classification

<sup>3</sup> Alteration of the production and secretion of sexual hormones in rats in some animal studies (ATSDR, 2017)

<sup>4</sup> Infertility in rats in some animal studies (ATSDR, 2017)

<sup>5</sup> There are also inorganic phosphorus-based flame retardants such as red-phosphorus and phosphates (Aschberger et al., 2017)

<sup>6</sup> Often associated with phosphorus-based flame retardants (US EPA, 2015)

Legend: BPA-BDPP, bisphenol A bis(diphenyl phosphate); CFCs, chlorofluorocarbons; DBDPE, decabromodiphenyl ethane or 1,2-bis(pentabromodiphenyl)ethane; DBHCTD; dibromo-hexachlorotricyclotridecene; DPs, dechlorane plus; DOPO, 9,10-dihydro-9-oxa-10-phosphaphenanthrene-10-oxide; HBCD, hexabromocyclododecane; HC, hydrocarbons; HCFC, hydrochlorofluorocarbons; HFC, hydrofluorocarbons; PBDPP, 1,3-phenylene bis(diphenyl phosphate) or resorcinol bis(diphenyl phosphate); PCBs, polychlorinated biphenyls; PFOS/F, Perfluorooctane sulfonate; PVC, polyvinyl chloride; TBBPA, tetrabromobisphenol A; TBBPA-BGE, tetrabromobisphenol A-bis bis(glycidyl)ether; TBPH, bis(2-ethylhexyl)-3,4,5,6-tetrabromo-phthalate; TCEP, tris(2-chloroethyl) phosphate; TCIPP, tris(2-chloroisopropyl) phosphate; TDCIPP, tris(1,3-dichloroisopropyl) phosphate; TPHP/TPP, triphenyl phosphate.  
- CRTs, cathode ray tubes; IT, information technology; LCDs, Liquid Crystal Displays; LED, light-emitting diode; PTVs, Plasma televisions; PCs, personal computers; TVs, televisions; WPCBs, waste printed circuit boards.

(Based on information provided by Cummings et al., 2012; Lundgren, 2012; Kiddee et al., 2013; Fornalczyk et al., 2013; Klaassen, 2013; Namias, 2013; Haque et al., 2014; Julander et al., 2014; Szalatkiewicz, 2014; Baldé et al., 2015; Bellanger et al., 2015; Cucchiella et al., 2015; 2016; Son et al., 2015; US EPA, 2015; Zheng et al., 2015; Cayumil et al., 2016; Gramatica et al., 2016; Matsukami et al., 2016 and 2017; Nartey, 2016; Someya et al., 2016; Woo et al., 2016; American Chemistry Council, 2017; ATSDR, 2017; Aschberger et al., 2017; IARC, 2017; Tansel, 2017; Toxnet, n.d.).

339 Human exposure to e-waste contaminants may be occupational or environmental, affecting both workers  
340 and the general population. As mentioned before, informal e-recycling sectors may involve not only adult  
341 workers but also children, teenagers and older adults, who are mainly involved in outdoor activities,

including dismantling and open burning. While most of formal e-recycling facilities have a particular concern for worker protection and are more able to invest in safe practices and disposal, workers involved in informal e-recycling activities are generally untrained and not equipped with personal protective equipment. They tend to use risky practices and sometimes work in worrisome conditions with no efforts to protect themselves, mainly due to the lack of knowledge of or access to protective measures (Iqbal et al., 2015; Adesokan et al., 2016; Amankwah-Amoah, 2016; Ceballos and Dong, 2016; Terazono et al., 2017).

Exposure may also occur through pollutants persisting in different environmental matrices such as air, surface water and groundwater, soil, sediment, food, and wastewater. As workers involved in the informal sector usually live within close vicinity of or even within recycling sites, they may be even more exposed to those contaminants. Exposure pathways are multiple, depending not only on the nature of the e-waste processing, but also on the type of contaminants involved. They may include inhalation, ingestion, dermal contact, transplacental and lactational routes (Lundgren, 2012; Grant et al, 2013; Kedee et al., 2013; Iqbal et al., 2015; Asampong et al., 2015; Carlson, 2016).

As is the case with other toxic waste, environmental damage from a lengthy history of inadequate e-waste management is expected to linger for a long time after the closure or upgrading of e-waste recycling sites further to the implementation of tighter environmental regulations or a shift from informal practices to more regulated ones (Xu Xinjin et al., 2015; Ceballos et al., 2016; Wang Yan et al., 2016). This is reflecting the e-waste legacy contamination, which is even more critical given the sustained involvement of persistent and bioaccumulative organic pollutants. According to LeBel (2016), exposure to e-waste

362 contaminants may “defy the temporal practices and qualities associated with traditional waste  
363 management strategies.”

364 **Table 2**

365 **Some secondary and tertiary e-waste contaminants released during e-waste processing**

E-waste processing	Most common contaminants emitted	Category of contaminants, examples of source or use	Main types of toxicity
Combustion of e-waste containing PCBs, plastics and PVC in order to recover precious materials (open burning activities, pyrometallurgical process in a furnace)	PAHs (e.g., phenanthrene, anthracene, fluoranthene, benzo [a] pyrene, benz [a] anthracene)	Secondary (from brominated, chlorinated and organophosphorus FRs in plastics and from PVC)	Carcinogenic <sup>1</sup> , photosensitizer
	PCDD/Fs		Immunotoxic, carcinogenic, reprotoxic, endocrine disrupters, may induce birth defects and also dermal damage (chloracne),
	PBDD/Fs		
	PXDD/Fs		
Incineration of e-waste residues as disposal strategy	Bisphenol A	Secondary (from combustion of polycarbonate plastics)	Endocrine disrupter
	Gases such as carbon monoxide and methane	Secondary (generated during smelting process)	Asphyxia
	Acids	Secondary (e.g., hydrobromic acid from brominated FRs, hydrochloric acid from chlorinated FRs and from incomplete combustion of PVC; phosphoric acid from organophosphorus FRs)	Induce from mild to severe burns to eyes and skin, sore throat, respiratory problems, corrosive injuries to lips, mouth, throat, etc., if swallowed.
Pyrometallurgy for metal refining	Acids (e.g., hydrochloric, nitric, sulphuric)	Tertiary (used as electrolyte solutions for electro-refining)	See above
Hydrometallurgy for metal refining	Acids (e.g., hydrochloric, nitric, sulphuric)	Tertiary (used as leaching agents, for solvent-extraction and for electroextraction solutions)	See above
	Cyanides	Tertiary (used as reagents)	Neurotoxic, cardiotoxic, may induce chest pain, breathing difficulties, increase in size of thyroid gland
	Thiourea	Tertiary (used as reagents)	Photosensitizer, hematotoxic,

<sup>1</sup> Some PAHs are *probably carcinogenic to humans* (e.g., ben[a]anthracene and benzo[a]pyrene), while some are *possibly carcinogenic to humans* (e.g., benzo[a]fluoranthene and benzo[k]fluoranthene), according to the IARC classification (ATSDR, 2016).

Legend: FRs, flame retardants; PBDD/Fs, polybrominated dibenzo-p-dioxins and dibenzofurans; PAHs, polycyclic aromatic hydrocarbons; PCBs, polychlorinated biphenyls; PCDD/Fs, polychlorinated dibenzo-p-dioxins and dibenzofurans; PVC, polyvinyl chloride; PXDD/Fs, mixed polybromochloro-dibenzo-p-dioxins and dibenzofurans.

(Based on information provided by Schluep et al, 2009; Lundgren et al., 2012; Grant et al., 2013; Kiddee, 2013; An et al., 2014; Cayumil et al., 2016; Chen Jiangyao et al., 2016; Zhang Tao et al., 2016; Aschberger et al., 2017; ATSDR, 2017; Iannicelli-Zubiani et al., 2017; IARC, 2017; Toxnet, n.d.)

#### 4.1. Legacy contamination and human body burden

The environmental burden associated with e-waste contaminants began receiving attention from environmental advocacy groups in the 1990s (Ahmad Khan, 2016). A 2002 report from two non-governmental organizations (NGOs) – the Basel Action Network and the Silicon Valley Toxics Coalition (SVTC) – described some particularly harmful e-recycling activities (e.g., open-pit acid baths, smelting operations on the ground, crushing and open dumping of CRT TVs along rivers, open burning of wires within cities and near rivers and streams) with the complete absence of personal protective measures in Guiyu (China), Sher Shah (Pakistan) and New Delhi (India). Water sampled from rivers adjacent to WPBCs burning sites in Guiyu revealed lead concentrations 190 times higher than those set in the World Health Organization (WHO) Drinking Water Guidelines in 2001. Sediments sampled from the same location showed barium, tin and chromium concentrations respectively 10, 152, and 1,338 times higher than the US EPA threshold for environmental risk in soil (BAN/SVTC, 2002).

Despite regulatory efforts and technological progress over the years, studies continue to uncover human exposure and environmental contamination attributable to e-waste handling and recycling activities, especially in developing countries (Lundgen, 2012; Song et al, 2015; Du et al., 2016; Awasthi et al., 2016). Table 1, in the Appendix, presents some of the most recent field studies (including a few human risk assessments) that have documented environmental impacts attributable to e-recycling practices. The targeted contaminants were mainly heavy metals and POPs (flame retardants, PCBs, PAHs). Children constitute the population most sensitive to e-waste exposure as they are likely to receive high doses of contaminants, in particular because of their exposure through multiple routes (e.g., breastfeeding, placental, dermal, hand-to-mouth, object-to-mouth, take-home exposure), their higher basal metabolism

than adults, their larger surface area in relation to body weight and their lower toxin elimination rates (Zeng Xiang et al., 2016).

Several studies have explored the body burden resulting from e-waste exposure, not only in formal and informal workers but also in the surrounding population. A literature review by Ceballos and Dong (2016) reported that, despite the use of advanced technologies, workers' exposure to some toxic metals in formal e-recycling facilities exceeds the occupational exposure limits while biological levels of flame retardants were found to be higher than those of unexposed reference groups. These authors suggested that engineering and administrative controls can lower occupational exposure, which means that protective measures implemented thus far in some licensed facilities have failed to create a completely safe work environment. Investigating occupational exposure in CRT recycling facilities in France, Lecler et al. (2015) identified a few highly polluting steps (e.g., CRT decontamination) and concluded that there is a need for general ventilation, better occupational risk management and medical biomonitoring. High exposure to heavy metals, furans, dioxins, PCBs and other POPs were mostly reported for workers in the informal sectors and the surrounding population (Song et al., 2015; Cayumil et al., 2016; Edwards, 2016; Li et al., 2017). Table 2, in the Appendix, reports on recent studies that have carried out personal air and/or biological sampling to investigate the body burden of e-waste exposure in workers and surrounding populations.

However, ongoing investigations on e-waste legacy contamination are revealing emerging issues that also raise concerns, thus complicating the situation even further. Some of these are detailed in Table 3.

### **Table 3**

#### **Emerging issues regarding the e-waste legacy contamination**



Emerging issues	Comments
Increased environmental and human burdens of a mix of typical and new persistent and bioaccumulative components, used as alternatives to PBDEs and/or plasticizers	<p>Accumulations of new flame retardants (e.g., bisphenol A bis(diphenyl phosphate) [BPA-BDPP], dechlorane plus [DPs], (2-ethylhexyl)diphenyl phosphate [EHDPP], hexabromobenzene [HBB], tris(2-chloroethyl) phosphate [TCEP], bis(1,3-dichloro-2-propyl) phosphate [TDCIPP], decabromodiphenyl ethane [DBDPE], triphenyl phosphate [TPHP]) found in dust, soil, sediment and food in informal e-recycling areas in China (Wu Ping-Fan et al., 2016; Zheng Xiaobo et al., 2016; He et al., 2017) and Vietnam (Someya et al., 2016; Matsukami et al., 2016 and 2017; Tao et al., 2016).</p> <p>High concentrations of organophosphate metabolites, deriving from PBDE alternatives (TCEP, TDCIPP, TPHP) also found in urine of people living and/or working in informal e-recycling areas of Longtang town, China (Lu et al., 2017).</p>
Environmental accumulation of emerging dioxin-related compounds	Accumulation of dioxin-like compounds in soil and sediment of informal e-recycling area in Vietnam. Toxic equivalent (TEQ) exceeding the Japan and Netherlands' criteria for human health (Suzuki et al., 2016). High levels of mixed halogenated (brominated and chlorinated) dibenzo-p-dioxins/dibenzofurans (PXDD/Fs) also found in soil of burning site in Agbogbloshie, Ghana (Tue et al., 2016b).
Release of potentially toxic volatile organic compounds (VOCs) during e-waste combustion	Waste printed circuit boards (WPCBs) and plastic casings' burning release high concentrations of VOCs, heavy metals and POPs. Aromatic hydrocarbons (AHs) (e.g., benzene, toluene, ethyltoluene, and phenol) identified as the dominant group of emitted VOCs, implying both non-cancer and cancer risk for workers, especially when using rotary incinerators. Predominance of AHs due to bisphenol A in the epoxy formulation in WPCBs (An et al., 2014; Chen Jiangyao et al., 2016). Combustion of WPCBs releases a complex mixture of AHs and aromatic amine hydrocarbons, including aniline (Sahle-Demessie et al., 2016).
Migratory effect and pollution scale	<p>Potentially high environmental migratory abilities of alternative flame retardants (particularly DPs and oligomeric organophosphorus [o-PFRs]) and their breakdown products, resulting in weathering processes during e-waste open storage and burning (Matsukami et al., 2017).</p> <p>Higher scale of pollution by organic pollutants (e.g., halogenated flame retardants, dioxin-like compounds, polyaromatic hydrocarbons [PAHs]) estimated, compared to metals, which may increase contamination risks for adjacent and surrounding areas and increase human health risk impacts (Fujimori and Takigami, 2014; Xiao et al., 2016; Terazono et al., 2017; Yuan et al., 2017). Outflow of PAHs from e-recycling activities in Guiyu (when facilities were in operation) may affect local population and also population living up to 30 km downstream in coastal region of Haimen Harbour (Shi et al., 2016).</p>
Other significant contributors to environmental burden	<p><b>Abandoned e-recycling sites</b></p> <p>Environmental persistence and potential dissemination of e-waste contaminants from abandoned e-recycling sites remain a human health risk years after closure of these sites. Abandoned burning sites in China remain a source of polychlorinated biphenyls (PCBs), especially during warmer seasons (Wang Yan et al., 2016; 2017). Heavy metals found to accumulate in ponds after being transported by rainfall from contaminated surface soil of abandoned e-recycling areas, thus posing a serious human health risk via dietary intake of agricultural products (Wu Qihang et al., 2015).</p>

	<p><b>Formal e-recycling sector</b> Soil contaminated with heavy metals, PCBs, brominated flame retardants and dioxin-related compounds (DRCs) within and around a e-recycling compound in the southeastern part of Tianjin City (Zhao et al., 2013; Hong et al., 2016). Same results reported for an emerging large-scale complex in Taizhou (Ma et al., 2008; 2009).</p> <p>Several officially licensed facilities, which are engaged in low-tech e-recycling practices, are still carrying out rudimentary activities such as open burning of wires and cables (Ma et al., 2008 and 2009; Hong et al., 2016).</p> <p>There have been instances of formal facilities in the US involved in legal litigation for leaving contaminated sites after bankruptcy or for illegally dumping e-waste on abandoned lots (Elliot, 2016).</p>
Other influencing factors to consider in human health risk assessment	<p>Migrant e-recycling workers in Taizhou, China, had PCB-related health risk 3.8 times greater than that of local residents, both for cancer and non-cancer risks. Suggestion made that non-dietary exposure (e.g., lifestyle and economic status) should also be considered in human health risk assessment among workers (Wang Yalin et al., 2016).</p> <p>In Agbogbloshie, Ghana, lower exposure to PCBs found for e-recycling workers. Low living standard, associated with less fish consumption, the major dietary source of PCB exposure (Wittsiepe et al., 2015).</p>
Other critical exposure routes	<p>Dermal uptake from e-waste airborne contaminants (skin absorption of particulates and gaseous organic contaminants from e-waste combustion fumes) considered a significant exposure route for e-recycling workers and populations living in and around recycling areas, especially for organic pollutants such as flame retardants and plasticizers with low molecular weights that allow high dermal penetration rates and large skin-air partition coefficients (Wu Chenchou et al., 2016).</p>

## 4.2. Remaining gaps in scientific knowledge in legacy contamination

Despite scientific interest in the e-waste legacy contamination, many knowledge gaps remain regarding its complexity and the scale of its potential environmental damage, but also in the identification of adverse health effects attributable to e-waste contaminants. Table 4 presents some important scientific knowledge gaps that still need to be bridged for a better assessment of the human and environmental burdens of e-waste management. The most significant gap is probably the lack of direct evidence of causal relationships between exposure to e-waste contaminants and adverse health effects, in part due to inconsistencies in epidemiological data, absence of dose-response relations and still-poor documentation of biological

416 mechanisms of the effects of e-waste contaminant mixtures (Grant et al., 2013; Xu Xijin et al., 2015; Zeng  
417 Xiang et al., 2016).

#### 418 **Table 4**

#### 419 **Some scientific knowledge gaps regarding the e-waste legacy contamination**

<b>Knowledge gaps</b>	<b>Comments</b>
Comprehensive characterization of e-waste contaminants including specific properties (e.g., toxicity, environmental behaviour, potential interaction between contaminants in the environment)	<ul style="list-style-type: none"> <li>• Unidentified halogenated PAHs are released during e-waste combustion. Their potential toxicity is still unknown to date (Tue et al., 2016a).</li> <li>• Speciation and leaching behaviour of metals in e-waste contaminated areas have been little studied (Cui et al. 2017).</li> <li>• More studies are needed to assess trophic transfer of hexabromocyclododecanes (HBCD) in the food webs (Zhu et al., 2017).</li> <li>• There are still uncertainties about alternative flame retardants with regard to their environmental behaviour, potential toxicity, metabolism, persistence and potential of bioaccumulation in biota (Matsukami et al., 2016 and 2017; Tao et al., 2016; Zhu et al., 2017).</li> <li>• The interplay between contaminants is still poorly documented; however, Pb, Cu, Zn, and Br contained in contaminated soil and ash mixture of open e-waste burning areas in Agbogbloshie, Ghana, are thought to influence the formation of dioxin-related compounds (Fujimori et al., 2016).</li> </ul>
Characterization of human exposure to alternative flame retardants	<ul style="list-style-type: none"> <li>• More studies are needed to investigate human exposure to alternative flame retardants and assess their human health risk (Matsukami et al., 2017). Levels of some novel brominated flame retardants in samples of foodstuffs exceeded those in foods from control sites in informal e-recycling area in Taizhou City, Eastern China (Labunska et al., 2015).</li> </ul>
Human health exposure and risk assessment related to multiple exposure from e-recycling	<ul style="list-style-type: none"> <li>• Human health risks related to multiple exposure in developing countries, for workers and surrounding populations, are still poorly documented (Grant et al., 2013; Ceballos and Dong, 2016; Xu Long et al., 2016; Zeng Xiang et al., 2016; Akortia et al., 2017). Despite enduring exposure of Ghanaian e-recycling workers to the highest levels of chlorinated and brominated dioxin-like compounds ever reported, cancer risks related to complex mixtures of potential mutagenic PAHs and derivatives are also still poorly assessed (Tue et al., 2016a).</li> <li>• Data and awareness on take-home exposure are still lacking (Perkins et al., 2014; Ceballos and Dong, 2016). Children are those mostly confronted with take-home contamination in both developing and developed countries (Zeng Xiang et al., 2016).</li> <li>• To date, little research has focused on workers' exposure or risk assessment in formal e-recycling. More studies are required to assess progress in occupational exposure following implementation of better protective measures in formal e-recycling and to evaluate improvement in environmental and human health exposure following a shift from informal activities to more formal ones in developing countries (Ceballos and Dong, 2016).</li> <li>• Neither exposure assessment nor health risk assessment has been conducted among surrounding populations of formal e-recycling facilities in developed countries.</li> </ul>

Knowledge gaps	Comments
Understanding of the toxicology of chemical mixtures in e-recycling	<ul style="list-style-type: none"> <li>• More studies are required to investigate potential inhibitory, synergistic, or additive effects following multiple exposures to e-waste components in order to adequately assess human health risks (Grant et al., 2013). For example, experimental repeated exposure of earthworms to BDE-209/Pb (bromodiphenyl ether congener/lead) appears to increase the production of reactive oxygen species (ROS) (Hu et al., 2016). Another experiment led to contradictory results when assessing the interaction of Cd and flame retardant BDE-209 on blood cells with different evaluation methods (Curcic et al., 2017).</li> </ul>
Better characterization of legacy contamination in developing countries	<ul style="list-style-type: none"> <li>• China remains the focus of scientific interest due to its long informal e-recycling history and its position as a leading hub in the e-waste trade (Lines et al., 2016). However, there is growing concern about the prevalence of brominated flame retardants in various environmental matrices in other Asian countries (Mackintosh et al., 2015).</li> <li>• Few field studies of environmental issues related to informal e-recycling have been conducted in India, which is surprising as it plays a non-negligible role in the current imported e-waste trade, after China (Borthakur and Singh, 2017).</li> <li>• Ghana has been a focus of scientists with regard to its heavy contamination legacy, but the situation in other African countries, such as Nigeria, remains poorly documented. However, this country was involved in the hazardous waste trade early on and is becoming a hub for the second-hand market, while large open dumpsites are proliferating virtually everywhere (Peluola, 2016).</li> </ul>
Characterization of legacy contamination related to e-waste landfilling and formal e-recycling in developed countries	<ul style="list-style-type: none"> <li>• In developed countries, very few field studies similar to the work of Klees et al. (2017) (see Table 1 in Appendix A) have been carried out to date to address potential releases of e-waste contaminants from formal e-recycling facilities or landfill sites and their accumulation in the surrounding environment, despite calls for such research from a few authors (Ceballos and Dong, 2016; Seeberger et al., 2016). A human health and ecological risk assessment for e-recycling facilities conducted in Canada in 2004 had already underlined the need for further research on occupational health hazards related to e-waste contaminants, and for further environmental monitoring to assess potential risks for the surrounding population and environment (MJC &amp; Associates, 2004). It is worth to mention that the increase in landfilling of CRT is already happening in the US as a response to the difficulty in processing the large quantities of CRT waste (Elliot, 2017).</li> </ul>
Stronger body of evidence of adverse human health effects specifically ascribed to e-waste contaminants	<ul style="list-style-type: none"> <li>• Cohort studies with sufficient sample sizes are required to explore the causal relationship between exposures to e-waste and detrimental health effects, especially among children and pregnant women subject to long-term exposure to e-waste contaminants. The existing epidemiological studies are descriptive and plagued by small numbers, sampling bias and inadequate adjustment for confounding factors (Grant et al., 2013; Xu Xinjin et al., 2015; Lin et al., 2016).</li> <li>• No health studies have been done so far on formal e-recycling workers, in both developed and developing countries (Ceballos and Dong, 2016), or on populations surrounding formal e-recycling facilities and landfill sites in developed countries.</li> </ul>
More effective land decontamination technologies	<ul style="list-style-type: none"> <li>• Remediation of e-waste contaminated lands is poorly documented, as field studies and full-scale applications are still overlooked (Duan et al., 2016). Available technologies are still at the experimental stage, including remediation of PCB-contaminated soils using plants (e.g., rice, alfalfa, ryegrass) (Shen et al., 2009), ultrasound-assisted soil washing to remove POPs and heavy metals (Chen Fu et al., 2016), and bioremediation techniques using bacteria with the ability to absorb lead present in contaminated soil (Gayatri and Shailaja, 2016).</li> </ul>

Knowledge gaps	Comments
Contribution of social factors in the increase of exposure and negative health outcomes among e-recycling workers.	<ul style="list-style-type: none"> <li>• Some workers face increased risk for occupational injuries and illnesses as a result of social and economic structures historically linked to discrimination or exclusion. These are known as occupational health inequalities (Leong et al., 2017).</li> <li>• By in large, e-waste recycling is conducted by workers from social marginalized groups such as prisoners, immigrants, ethnic minorities, and children.</li> <li>• More research is needed to explore how social institutions such as racism, xenophobia, sexism, and classism materialize in the e-waste recycling industry and contribute to an increased risk of negative health outcomes for certain workers or segments of the general population (i.e. environmental justice) (NIOSH/ASSE, 2015).</li> <li>• More research needs to be conducted on how the structure of the e-waste industry puts workers at greater risk for negative health outcomes. Topics of interest include how factors such as employment arrangements such as temporary employment or prison work impact worker safety.</li> <li>• Similarly, the relationship between organizational factors - such as business size and formal/informal sectors – and worker health should be explored.</li> </ul>

## 5. Potential solutions to curb e-waste issues

Solving e-waste issues is a long-term, challenging and costly task and calls for the full collaboration of all stakeholders involved (e.g., government, scientific community, chemical manufacturers, e-product designers, retailers and collectors, e-recyclers and end consumers). However, any action taken to address e-waste problems must be strongly backed by a firm governmental position in terms of tighter laws, regulations and policies, including surveillance of the Extended Producer Responsibility (EPR) system, awareness initiatives, sustained financial, technical, logistical, and training support, and even international cooperation. The aim should be to steadily increase the engagement and adhesion of other stakeholders (Gu et al., 2016; Heacock et al., 2016; Step, 2016; Zhang and Guan, 2016; Zeng et al., 2017). That being said, the gaps have widened between rapidly evolving technological markets, high consumption of e-products and, hence, high generation of e-waste, on the one hand, and the resources available to deal adequately with the potentially hazardous e-waste flow, on the other. For instance, despite three decades of intense research, the performance and stability of substitute materials for e-product manufacturing are still problematic (Li et al., 2015). Yet, given the serious human health risks and environmental burden, it

is imperative that the e-waste problem-solving process gain momentum. A comprehensive approach is suggested here that involves waging a battle on two fronts: at the upstream and downstream levels.

Potential upstream solutions could include:

- **Substantially reducing the use of potentially toxic compounds.** Some substitutes for hazardous e-waste components already exist (e.g., lead-free alloys to replace traditional tin-lead solder, a nanoplate cobalt alloy instead of Cr, and carbon nanotubes to replace metal particles). However, a few substances such as Cd, Hg, Cr and flame retardants remain more challenging to replace in terms of physical and chemical properties. Furthermore, the affordability and functionality of safer materials remain the primary concern for e-product designers (Li et al., 2015; Meyer and Katz, 2016; You et al., 2016). Technology convergence, such as the integration of multiple devices into a single versatile one, may be an effective way to reduce human health and environmental impacts, provided that new devices replace as many single ones as possible (Son et al., 2015). Another avenue might be packaging designed to induce consumers to choose more environmentally friendly products (e.g., short description of chemical composition on e-product packaging) (You et al., 2016). Also, green electronics, which are still at the experimental phase, require further research and evaluation, global regulations, better consumer awareness and stronger incentives (Verdecchia et al., 2017).
- **Decreasing quantities of e-waste.** To date, the main focuses have been on waste management and clean production. However, a promising new avenue is that of joint mobilization (including strong governmental involvement) against the functional and subjective obsolescence of e-products by endeavoring to increase both manufacturers' and consumers' accountability. A more

visible relationship needs to be established between planned and subjective obsolescence and potentially negative health effects and environmental degradation, on both the local and global scales. Profound changes in consumption patterns are also in order, without infringing on the individual's right to consume. Education and awareness-raising initiatives should be undertaken on a broader scale to reach large audiences, with sensitization beginning during formal education as early as elementary school. Achieving a higher level of awareness would empower consumers to become change drivers who could push manufacturers towards more sustainable business approaches (Wilson et al., 2015; Ahmed Khan, 2016; Echegaray, 2016). Manufacturers also have a role to play in influencing consumer behaviours through more ethical attitudes and a lasting commitment to sustainable production. One of manufacturers' main contributions could be to ensure sustainable designs and optimize the feasibility of e-product repairs, thus guaranteeing a welcome increase in the lifespan of e-devices. Other possible solutions could include more flexible assembly modes such as bolt connections instead of rivet connections, standardized motherboards with fixed connections, better technical supervision of repairers and greater visibility for repair service shops (O'Connor et al., 2016; You et al., 2016; Sabbaghi et al., 2017). Studies have also shown the potential for a technical increase in e-products' lifespan: manufacturers are therefore requested to encourage research and development in this promising area (Yadav et al., 2015). In this context, it is worth underlining the efforts of the European Parliament which adopted a motion on "*a longer lifetime for products*" in June 2017, for the first time calling on the European Commission to legislate against planned obsolescence (European Parliament, 2017).



- **Increasing e-waste recyclability.** Many strategies have been suggested to promote eco-design right from the conception stage by adding an environmental dimension to the existing resource, thus improving the feasibility and added value of e-recycling processes. This includes concepts such as design for environment (DFE), design for recycling (DFR), design for disassembly (DFA), and design for recovery (DFR). Intended to increase the ease of both repair and recycling processes, these strategies are mostly based on modular designs integrating dismantling process facilitators (e.g., fewer screws, use of snap fits, metal frames instead of plastic ones, separable touch screens and batteries) and on optimal material compatibility when selecting material combinations based on both their recycling properties and their performance criteria (Li et al., 2016; Movilla et al., 2016; O'Connor et al., 2016; Zeng et al., 2017).

Potential downstream solutions could include:

- **Stopping the illegal trade while upgrading the informal e-recycling sector, pending full implementation of a safer formal one.** These two actions go hand-in-hand, given that the illegal export of e-waste feeds informal practices. However, even if there is a total ban, an informal e-recycling sector will continue to handle stockpiled imported and domestic e-waste. This remains a very important means of subsistence for low-income and less-educated populations. Cracking down on this sector by reinforcing regulation would do more harm than good, as unemployment would rise and hence degrade living and health conditions among already vulnerable populations. A more pragmatic solution would be to provide a better framework to the informal sector through the increase of awareness, education, training and occupational and environmental health guidance among informal workers, along with tools and technical and financial support (Amankwaa et al.,



2016; Ardi and Leisten, 2016; Heacock et al., 2016). The “Best of two worlds” (Bo2W) model, developed by the multi-stakeholder Step Initiative, advocates a division of tasks such that a more protected informal sector in developing countries would take part only in manual dismantling activities (referred to as the “best” pre-processing), while the complex e-waste fraction loaded with hazardous and valuable materials would be safely treated in state-of-the-art facilities located in developed countries (referred to as the “best” end-processing). Manual dismantling is thus considered more efficient and less costly for a higher yield of material liberation than the mechanical option. The Bo2W philosophy is seen as a transition measure and even as a “global reverse logistic” until such time as high-tech e-recycling facilities are fully implemented in developing countries (McCann and Wittermann, 2015; Wilson et al., 2015; Geeraerts et al., 2016; Singh et al., 2016). Similarly, some authors also advocate a “*regulated green transboundary channel*” of e-waste between developed and developing countries to sustain the reuse market, in addition to dismantling activities, suggesting that exporting countries could also fund appropriate e-waste management in receiving countries (Milovantseva and Fitzpatrick, 2016). This action would be extremely relevant, given the quantity of still-operational devices stored at home in developed countries, as revealed by surveys conducted in Canada D  m  n   and Marchand, 2016; Dewis and Van Wesenbeeck, 2016) and in the United Kingdom (Wilson et al., 2017). International cooperation is also suggested to support formal e-recycling in developing countries, mostly in terms of technology transfer and certification of facilities complying with legal and environmental criteria (Zhang and Guan, 2016).

- 517       • **Introducing stronger and innovative reverse logistics to close the loop.** Reverse logistics  
518 encompasses the processes of recovering used goods. In the case of e-waste, it includes all the  
519 mechanisms implemented to encourage consumers to properly dispose of e-waste and return e-  
520 products to the manufacturers. Reverse logistics is a great concern for all countries. Advocated  
521 proactive actions include increasing consumers' knowledge and awareness of the human health  
522 and environmental risks associated with inappropriate practices, and boosting their willingness to  
523 be proactive (e.g., through participation in policy design and implementation, which includes more  
524 convenient modes of returning devices; and through multiplication of strategic collection points;  
525 environmentally oriented education to instill moral attitudes and behaviours; and use of social  
526 media and advertising campaigns). Other proactive actions could include the creation of more  
527 formalized jobs in collecting and sorting areas, together with the inclusion, education and training  
528 of informal workers, and the promotion of tax and credit incentives among manufacturers, retailers  
529 and formal collectors (Dixit et al., 2016; Grant and Oteng-Abadio, 2016; Gu et al., 2016; Guarnieri  
530 et al., 2016; Liang and Sharp, 2016). In developed countries, better cooperation between the private  
531 and public sectors could be more productive as competitive collective systems may play a  
532 significant role in improving collection rate from households than a single or non-competing  
533 collective system (Corsini et al., 2017).
- 534       • **Upgrading formal e-recycling processes.** Intensive research is currently under way to enhance  
535 e-recycling technologies and profitability in order to ensure optimal recovery of valuable  
536 materials and thus reduce the pressure on the demand for raw materials, to create sustainable  
537 businesses and jobs, to treat hazardous fractions in an environmentally sound manner, and

ultimately to dispose of e-waste properly (Kumari et al., 2016; Li et al., 2016; Movilla et al., 2016; Iannicelli-Zubiani et al., 2017; Tansel, 2017). Banning the export of e-waste would undoubtedly provide a strong incentive to improve e-recycling processes. In these times of scarce resources and environmental degradation (Ryneal, 2016; Tansel, 2017), upgrading formal e-recycling processes would be doubly advantageous, given the great economic development and profitability potential and the gainful employment opportunities they offer in developing countries (Amankwah-Amoah, 2016; Ryneal, 2016), as well as in developed countries (Cucchiella et al., 2016).

## 6. Conclusion

To the best of our knowledge, this is the first paper to adopt a holistic approach that exposes the extreme complexity of the issues, challenges and gaps (including human health concerns and environmental degradation) associated with the tricky e-waste question, and that suggests potential solutions, both upstream and downstream. The many initiatives launched thus far to advocate suitable management practices are facing disappointing realities, which are reflected in heavy legacy contamination and significant human exposure to e-waste contaminants in surrounding populations and in both formal and informal e-recycling. Exploring e-waste issues is akin to opening a Pandora's box, with constantly emerging e-waste legacy contamination issues regarding, for example, critical exposure pathways, unexpected sources and environmental fates of contaminants, as well as novel e-waste contaminants, all of which is further complicated by an increasingly intricate illegal e-waste trade.

However, despite scientific gaps, current knowledge of the threats posed by persistent and bioaccumulative pollutants is sufficient to raise awareness and concern among all stakeholders. Particular efforts should be expended to improve e-recycling technologies in terms of efficiency, affordability and environmental performance, given the great economic potential of e-recycling and its key role in preventing the incineration and/or landfilling of e-waste. As consumer habits and discipline remain essential to the resolution of any e-waste problem, joint efforts are needed to fight e-products' subjective and functional obsolescence through the environmentally oriented education of consumers, which in turn will help induce manufacturers to make ethical and sustained commitments.

Human health and environmental integrity can quite conceivably be preserved while safely and wisely managing e-waste resources. However, achieving this may require an ideological shift toward a perception of e-waste as a vital input that could be re-introduced into other processes. This would effectively keep the lid on Pandora's box.

## **Acknowledgments**

This research was supported by the Institut de recherche Robert-Sauvé en santé et en sécurité du travail (IRSST), Montréal, Québec, Canada (grant no. 2015-0083).

## References

- Adesokan, M. D., Adie, G.U., Osibanjo, O., 2016. Soil pollution by toxic metals near e-waste recycling operations in Ibadan, Nigeria. *J. Health Pollut.* 6 (11), 26-33.
- Ahmad Khan, S., 2016. E-products, e-waste and the Basel Convention: Regulatory challenges and impossibilities of international environmental law. *RECIEL*. 25(2), 248-260.
- Akortia, E., Olukunle, O. I., Daso, A. P., Okonkwo, J. O., 2017. Soil concentrations of polybrominated diphenyl ethers and trace metals from an electronic waste dump site in the Greater Accra Region, Ghana: Implications for human exposure. *Ecotox. Environ. Safe.* 137, 247-255.
- Amankwaa, E. F., Bowman, J., Tsikudo, K. A., 2016. Recyclers at risk? Analysis of e-waste livelihoods and blood lead levels at Ghana's recycling hub, Agbogbloshie. Policy Brief 33113, June 2016, International Growth Center. (<http://www.theigc.org/wp-content/uploads/2016/07/Amankwaa-Tsikudo-2016-Policy-brief.pdf>, accessed 10.01.2017)
- Amankwah-Amoah, J., 2016. Global business and emerging economies: Towards a new perspective on the effects of e-waste. *Technol. Forecast. Soc.* 105, 20-26.
- American Chemistry Council, 2017. Classes of Flame Retardants: North American Flame Retardant Alliance. (<https://flameretardants.americanchemistry.com/Classes-of-FRs>, accessed 12.05.2017)
- An, T., Huang, Y., Li, G., He, Z., Chen, J., Zhang, C., 2014. Pollution profiles and health risk assessment of VOCs emitted during e-waste dismantling processes associated with different dismantling methods. *Environ. Int.* 73, 186-194.
- Anh, H. Q., Nam, V. D., Tri, T. M., Ha, N. M., Ngoc, N. T., Mai, P. T. N., et al., 2016. Polybrominated diphenyl ethers in plastic products, indoor dust, sediment and fish from informal e-waste recycling sites in Vietnam: a comprehensive assessment of contamination, accumulation pattern, emissions, and human exposure. *Environ. Geochem. Health.* 1-20. DOI: 10.1007/s10653-016-9865-6

595 Andrews, A., 2009. Beyond The Ban – Can The Basel Convention Adequately Safeguard the Interests of  
 596 the World's Poor in the International Trade of Hazardous Waste? LEAD Journal, 5/2, 167.  
 597 (<http://www.lead-journal.org/content/09167.pdf> [accessed 12.05.2017])

598 Ardi, R., Leisten, R., 2016. Assessing the role of informal sector in WEEE management systems: A  
 599 System Dynamics approach. Waste Manage. 57, 3-16.

600 Asampong, E., Dwuma-Badu, K., Stephens, J., Srigboh, R., Neitzel, R., Basu, N., Fobil, J. N., 2015.  
 601 Health seeking behaviours among electronic waste workers in Ghana. BMC Public Health, 15, 1065.

602 Aschberger, K., Campia, I., Pesudo, L. Q., Radovnikovic, A., Reina, V., 2017. Chemical alternatives  
 603 assessment of different flame retardants – A case study including multi-walled carbon nanotubes as  
 604 synergist. Environ. Int. 101, 27-45.

605 ATSDR (Agency for Toxic Substances & Disease Registry), 2017. ATSDR Toxic Substances Portal. Last  
 606 updated March 23, 2017. (<https://www.atsdr.cdc.gov/toxfaqs/index.asp>, accessed 26.05.2017)

607 Awasthi, A. K., Zeng, X., Li, J., 2016. Environmental pollution of electronic waste recycling in India: A  
 608 critical review. Environ. Pollut. 211, 259-270.

609 Baldé, C. P., Wang, F., Kuehr, R., Huisman, J., 2015. Global e-Waste Monitor – 2014. United Nations  
 610 University, IAS – SCYCLE, Bonn, Germany. ([https://i.unu.edu/media/ias.unu.edu-en/news/7916/Global-](https://i.unu.edu/media/ias.unu.edu-en/news/7916/Global-E-waste-Monitor-2014-small.pdf)  
 611 [E-waste-Monitor-2014-small.pdf](https://i.unu.edu/media/ias.unu.edu-en/news/7916/Global-E-waste-Monitor-2014-small.pdf) [accessed 10.01.2017]).

612 BAN/SVTC (Basel Action Network/Silicon Valley Toxics Coalition), 2002. Exporting Harm, the High-  
 613 Tech Trashing of Asia.  
 614 ([https://static1.squarespace.com/static/558f1c27e4b0927589e0edad/t/56184364e4b0cf5700f852af/14444](https://static1.squarespace.com/static/558f1c27e4b0927589e0edad/t/56184364e4b0cf5700f852af/1444430692735/Exporting+Harm+Report.pdf)  
 615 [30692735/Exporting+Harm+Report.pdf](https://static1.squarespace.com/static/558f1c27e4b0927589e0edad/t/56184364e4b0cf5700f852af/1444430692735/Exporting+Harm+Report.pdf), accessed 15.05.2017).

616 Basel Convention, 2011. Reports and Decisions. Decision IV/9 Adopted by the Conference of the Parties  
 617 at its Fourth Meeting, 6 May 1998,  
 618 (<http://basel.int/TheConvention/ConferenceoftheParties/ReportsandDecisions/tabid/3303/Default.aspx>,  
 619 accessed 10.01.2017).

Basel Convention, 2017. Amendment to the Basel Convention on the Control of Transboundary Movements of Hazardous Wastes and their Disposal, Official website. (<http://www.basel.int/Countries/StatusofRatifications/BanAmendment/tabid/1344/Default.aspx>, accessed 02.05.2017).

Bellanger, M., Demeneix, B., Grandjean, P., Zoeller, R. T., Trasande, L., 2015. Neurobehavioral deficits, diseases, and associated costs of exposure to endocrine-disrupting chemicals in the European Union. *J. Clin. Endocrinol. Metab.* 100(4), 1256-1266.

Borthakur, A., Govind, M., 2017. Emerging trends in consumers' E-waste disposal behaviour and awareness: A worldwide overview with special focus on India. *Resour. Conserv. Recy.* 117, Part B, 102-113.

Borthakur, A., Singh, P., 2017. Researches on informal e-waste recycling sector: It's time for a 'Lab to Land' approach. *J. Hazard. Mater.* 323 (Part B), 730-732.

Bradford, M., 2011. The United States, China & the Basel Convention on the Transboundary Movements of Hazardous Wastes and Their Disposal, *Fordham Environ. Law Rev.* 8(2), 305-349. (<http://ir.lawnet.fordham.edu/cgi/viewcontent.cgi?article=1445&context=elr>, accessed 26.05.2017).

Carlson, K., 2016. Electronic waste worker health: Studying complex problems in the real world. In: *Proceeding from the Eighth International Conference on Information and Communication Technologies and Development (ICTD)*, Article No. 35, Ann Arbor, Michigan, USA [June 3-6], ACM 978-1-4503-4306-0/16/06.

Cayumil, R., Khanna, R., Rajarao, R., Ikram-ul-Haq, M., Mukherjee, P. S., Sahajwalla, V., 2016. Environmental impact of processing electronic waste – Key issues and challenges. In: Mihai F.C. (Ed.), *E-waste in Transition - From Pollution to Resource*, InTech, Croatia, pp. 9-35.

Ceballos, D. M., Dong, Z., 2016. The formal electronic recycling industry: Challenges and opportunities in occupational and environmental health research. *Environ. Int.*

644 Ceballos, D., Beaucham, C., Page, E., 2017. Metal exposure at three U.S. electronic scrap recycling  
645 facilities. *J. Occup. Environ. Hyg.* 14(6), 401-408.

646 Chakraborty, P., Prithiviraj, B., Selvaraj, S., Kumar, B., 2016. Polychlorinated biphenyls in settled dust  
647 from informal electronic waste recycling workshops and nearby highways in urban centers and suburban  
648 industrial roadsides of Chennai city, India: Levels, congener profiles and exposure assessment. *Sci. Total*  
649 *Environ.* 573, 1413-1421.

650 Chen, A., Dietrich, K. N., Huo, X., Ho, S. M., 2011. Developmental neurotoxicants in e-waste: an  
651 emerging health concern. *Environ. Health Perspect.* 119(4), 431-438.

652 Chen, Fu, Yang, B., Ma, J., Qu, J., Liu, G., 2016. Decontamination of electronic waste-polluted soil by  
653 ultrasound-assisted soil washing. *Environ. Sci. Pollut. R.* 23(20), 20331-20340.

654 Chen, K., Zheng, J., Yan, X., Yu, L., Luo, X., Peng, X., et al., 2015. Dechlorane Plus in paired hair and  
655 serum samples from e-waste workers: correlation and differences. *Chemosphere.* 123, 43-47.

656 Chen, Jiangyao, Huang, Y., Li, G., An, T., Hu, Y., Li, Y., 2016. VOCs elimination and health risk  
657 reduction in e-waste dismantling workshop using integrated techniques of electrostatic precipitation with  
658 advanced oxidation technologies. *J. Hazard. Mater.* 302, 395-403.

659 Corsini, F., Rizzi, F., Frey, M., 2017. Extended producer responsibility: The impact of organizational  
660 dimensions on WEEE collection from households. *Waste Manag.* 59, 23-29.

661 Cucchiella, F., D'Adamo, I., Lenny Koh, S. C., Rosa, P., 2015. Recycling of WEEEs: An economic  
662 assessment of present and future e-waste streams. *Renew. Sust. Energ. Rev.* 51, 263-272.

663 Cucchiella, F., D'Adamo, I., Lenny Koh, S. C., Rosa, P., 2016. A profitability assessment of European  
664 recycling processes treating printed circuit boards from waste electrical and electronic equipments.  
665 *Renew. Sust. Energ. Rev.* 64, 749-760.

666 Cui, J. L., Luo, C. L., Tang, C. W. Y., Chan, T. S., Li, X. D., 2017. Speciation and leaching of trace metal  
667 contaminants from e-waste contaminated soils. *J. Hazard. Mater.*



668 Cummings, K.J., Nakano, M., Omae, K., Takeuchi, K., Chonan, T., Xiao, L., et al., 2012. Indium lung  
669 disease. *Chest*. 141 (6), 1512-1521.

670 Curcic, M., Buha, A., Stankovic, S., Milovanovic, V., Bulat, Z., Đukić-Ćosić, D., et al., 2017. Interactions  
671 between cadmium and decabrominated diphenyl ether on blood cells count in rats—Multiple factorial  
672 regression analysis. *Toxicology*, 376, 120-125.

673 D’Adamo, I., Paolo, R., Terzi, S., 2016. Challenges in waste electrical and electronic equipment  
674 management: a profitability assessment in three European countries. *Sustainability*, 8(633).

675 Damrongsiri, S., Vassanadumrongdee, S., Tanwattana, P., 2016. Heavy metal contamination characteristic  
676 of soil in WEEE (waste electrical and electronic equipment) dismantling community: a case study of  
677 Bangkok, Thailand. *Environ. Sci. Pollut. Res.* 23(17), 17026-17034.

678 Daso, A. P., Akortia, E., Okonkwo, J. O., 2016. Concentration profiles, source apportionment and risk  
679 assessment of polycyclic aromatic hydrocarbons (PAHs) in dumpsite soils from Agbogbloshie e-waste  
680 dismantling site, Accra, Ghana. *Environ Sci. Pollut. Res.* 23, 10883–10894.

681 Déméné, C., Marchand, A., 2016. Exploring users’ practices through the use phase of a television to  
682 minimise the environmental impact. *J. Res. Consumers.* 29, 4-13.  
683 ([http://www.jrconsumers.com/Consumer\\_Articles/issue\\_29/](http://www.jrconsumers.com/Consumer_Articles/issue_29/), accessed 10.01.2017).

684 Dewis G, Van Wesenbeeck, P., 2016. Trash talking - Dealing with Canadian household e-waste  
685 EnviroStats: Statistics Canada. Catalogue 16-002-X. ([http://www.statcan.gc.ca/pub/16-002-](http://www.statcan.gc.ca/pub/16-002-x/2016001/article/14570-eng.pdf)  
686 [x/2016001/article/14570-eng.pdf](http://www.statcan.gc.ca/pub/16-002-x/2016001/article/14570-eng.pdf), accessed 30.05.2017).

687 Dixit, S., Badgaiyan, A. J., 2016. Towards improved understanding of reverse logistics – Examining  
688 mediating role of return intention. *Resour. Conserv. Recy.* 107, 115–128.

689 Du, H., Yu, M., Sun, J., Song, G., Li, Y., 2016. Epidemiological evidence of the effects of environmental  
690 pollution on male reproductive health in an electronic waste-recycling town. *Pol. J. Environ. Stud.* 25(4),  
691 1777-1780.

692 Duan, H., Hu, J., Tan, Q., Liu, L., Wang, Y., Li, J., 2016. Systematic characterization of generation and  
 693 management of e-waste in China. *Environ. Sci. Pollut. Res. Int.* 23(2), 1929-1943.

694 Echegaray, F., 2016. Consumers' reactions to product obsolescence in emerging markets: the case of  
 695 Brazil. *J. Clean. Prod.* 134, 191-203.

696 Echegaray, F., Hansstein, F. V., 2017. Assessing the intention-behavior gap in electronic waste recycling:  
 697 the case of Brazil. *J. Clean. Prod.* 142, 180-190.

698 Edwards, L., 2016. E-Waste Recycling: The Dirty Trade between the United States and China. Pomona  
 699 Senior Theses. Paper 153. ([http://scholarship.claremont.edu/pomona\\_theses/153](http://scholarship.claremont.edu/pomona_theses/153), accessed 12.01.2017).

700 Eguchi, A., Nomiyama, K., Minh Tue, N., Trang, P. T., Hung Viet, P., Takahashi, S., Tanabe, S., 2015.  
 701 Residue profiles of organohalogen compounds in human serum from e-waste recycling sites in North  
 702 Vietnam: Association with thyroid hormone levels. *Environ. Res.* 137, 440-449. DOI:  
 703 10.1016/j.envres.2015.01.007

704 Elliot, B., 2016. Bankrupt processor leaves behind CRTs, E-Scrap News, Posted on March 10, 2016.  
 705 (<https://resource-recycling.com/e-scrap/2016/03/10/bankrupt-processor-leaves-behind-crts/>, accessed  
 706 06.08.2017)

707 Elliot, B., 2017. Landfilling CRT Glass Becomes the Norm in California. E-scrap News, Posted on March  
 708 2, 2017 ([https://resource-recycling.com/e-scrap/2017/03/02/landfilling-crt-glass-becomes-norm-](https://resource-recycling.com/e-scrap/2017/03/02/landfilling-crt-glass-becomes-norm-california/)  
 709 [california/](https://resource-recycling.com/e-scrap/2017/03/02/landfilling-crt-glass-becomes-norm-california/), accessed 06.08.2017).

710 European Commission, 2017. Waste Electrical & Electronic Equipment (WEEE). Last updated  
 711 2017/04/24. ([http://ec.europa.eu/environment/waste/rohs\\_eee/legis\\_en.htm](http://ec.europa.eu/environment/waste/rohs_eee/legis_en.htm), accessed 12.01.2017).

712 European Parliament, 2017. Report on a Longer Lifetime for Products: Benefits for Consumers and  
 713 Companies (2016/2272(INI)). Committee on the Internal Market and Consumer Protection, 9 June 2017  
 714 ([http://www.europarl.europa.eu/sides/getDoc.do?pubRef=-//EP//TEXT+REPORT+A8-2017-](http://www.europarl.europa.eu/sides/getDoc.do?pubRef=-//EP//TEXT+REPORT+A8-2017-0214+0+DOC+XML+V0//EN#title2)  
 715 [0214+0+DOC+XML+V0//EN#title2](http://www.europarl.europa.eu/sides/getDoc.do?pubRef=-//EP//TEXT+REPORT+A8-2017-0214+0+DOC+XML+V0//EN#title2), accessed 24.07.2017).

716 Flynn, M.A., 2014. Safety and the diverse workforce: Lessons from NIOSH's work with Latino  
717 immigrants. *Prof. Saf.* 59(6), 52-57.

718 Fornalczyk, A., Willner, J., Francuz, K., Cebulski, J., 2013. E-waste as a source of valuable metals. *Arch.*  
719 *Mater. Sci. Eng.* 63(2), 87-92.

720 Fujimori, T., Takigami, H., Agusa, T., Eguchi, A., Bekki, K., Yoshida, A., et al., 2012. Impact of metals  
721 in surface matrices from formal and informal electronic-waste recycling around Metro Manila, the  
722 Philippines, and intra-Asian comparison. *J. Hazard. Mater.* 221–222, 139-146.

723 Fujimori, T., Takigami, H., 2014. Pollution distribution of heavy metals in surface soil at an informal  
724 electronic-waste recycling site (Philippines). *Environ. Geochem. Health.* 36, 159-168.

725 Fujimori, T., Itai, T., Goto, A., Asante, K. A., Otsuka, M., Takahashi, S., Tanabe, S., 2016. Interplay of  
726 metals and bromine with dioxin-related compounds concentrated in e-waste open burning soil from  
727 Agbogbloshie in Accra, Ghana. *Environ. Pollut.* 209, 155-163.

728 Gayatri, Y., Shailaja Raj, M., 2016. Molecular characterization of bacteria isolated from e-waste dumping  
729 yards at Hyderabad, Telangana, India. *Eur. J. of Ex. Biol.* 6, 6:1. (<https://www.imedpub.com>, accessed  
730 12.01.2017).

731 Geeraerts K, Illes, A. I., Schweizer, J. P., 2016. Illegal Shipment of E-Waste from the EU: A Case Study  
732 on Illegal E-Waste Export from the EU to China. A Study Compiled as Part of the EFFACE project.  
733 London: IEEP, Institute for European Environmental Policy.  
734 ([http://efface.eu/sites/default/files/EFFACE\\_Illegal%20shipment%20of%20e%20waste%20from%20the](http://efface.eu/sites/default/files/EFFACE_Illegal%20shipment%20of%20e%20waste%20from%20the%20EU.pdf)  
735 [%20EU.pdf](http://efface.eu/sites/default/files/EFFACE_Illegal%20shipment%20of%20e%20waste%20from%20the%20EU.pdf), accessed 30.05.2017).

736 Ghosh, S. K., Debnath, B., Baidya, R., De, D., Li, J., Ghosh, S. K, et al., 2016. Waste electrical and  
737 electronic equipment management and Basel Convention compliance in Brazil, Russia, India, China and  
738 South Africa (BRICS) nations. *Waste. Manage. Res.* 34(8), 693-707.

739 Gramatica, P., Cassani, S., Sangion, A., 2016. Are some “safer alternatives” hazardous as PBTs? The case  
740 study of new flame retardants. *J. HazardL Mater.* 306, 237-246.

741 Grant, K., Goldizen, F. C., Sly, P. D., Brune, M.-N., Neira, M., van den Berg, M., Norman, R. E., 2013.  
 742 Health consequences of exposure to e-waste: a systematic review. *Lancet Glob. Health.* 1(6), e350-e361.

743 Grant, R. J., Oteng-Abadio, M., 2016. The Global Transformation of Materials and the Emergence of  
 744 Informal Urban Mining in Accra, Ghana. *Africa Today.* 62(4), 2-20. Published by Indiana University  
 745 Press. (<https://muse.jhu.edu/article/620263> , accessed 30.05.2017).

746 Gu, Y., Wu, Y., Xu, M., Wang, H., Zuo, T., 2016. The stability and profitability of the informal WEEE  
 747 collector in developing countries: A case study of China. *Resour, Conserv. Recy.* 107, 18-26.

748 Guarnieri, P., e Silva, L. C., Levino, N. A., 2016. Analysis of electronic waste reverse logistics decisions  
 749 using strategic options development analysis methodology: A Brazilian case. *J. Clean. Prod.* 133, 1105-  
 750 1117.

751 Guna, V. K., Murugesan, G., Basavarajaiah, B. H., Ilangoan, M., Olivera, S., Krishna, V., Reddy, N.,  
 752 2016. Plant-based completely biodegradable printed circuit boards. *IEEE Transactions on Electron*  
 753 *Devices.* 63(12), 4893-4898.

754 Guo, X., Yan, K., 2017. Estimation of obsolete cellular phones generation: A case study of China. *Sci.*  
 755 *Total Environ.* 575, 321-329.

756 Haque, N., Hughes, A., Lim, S., Vernon, C., 2014. Rare earth elements: Overview of mining, mineralogy,  
 757 uses, sustainability and environmental impact. *Resources.* 3, 614-635.

758 He, C. T., Zheng, X. B., Yan, X., Zheng, J., Wang, M. H., Tan, X., et al., 2017. Organic contaminants and  
 759 heavy metals in indoor dust from e-waste recycling, rural, and urban areas in South China: Spatial  
 760 characteristics and implications for human exposure. *Ecotox. Environ. Safe.* 140, 109-115.

761 Heacock, M., Kelly, C. B., Suk, W. A., 2016. E-waste: the growing global problem and next steps. *Rev.*  
 762 *Environ. Health.* 31(1), 131-135.

763 Honda, S., Khatriwal, D. S., Kuehr, R., 2016. Regional E-waste Monitor: East and Southeast Asia. United  
 764 Nations University ViE – SCYCLE, Bonn, Germany. ([http://ewastemonitor.info/pdf/Regional-E-Waste-](http://ewastemonitor.info/pdf/Regional-E-Waste-Monitor.pdf)  
 765 [Monitor.pdf](http://ewastemonitor.info/pdf/Regional-E-Waste-Monitor.pdf), accessed 30.05.2017).

766 Hong, W. J., Jia, H., Ding, Y., Li, W. L., Li, Y. F., 2016. Polychlorinated biphenyls (PCBs) and  
 767 halogenated flame retardants (HFRs) in multi-matrices from an electronic waste (e-waste) recycling site  
 768 in Northern China. *J. Mater. Cycles and Waste Manag.* 1-11.

769 Hopson, E., Puckett, J., 2016. Scam Recycling: e-Dumping on Asia by US Recyclers. The e-Trash  
 770 Transparency Project, Basel Action Network. ([https://www.resource-](https://www.resource-recycling.com/images/BANReportTwo.pdf)  
 771 [recycling.com/images/BANReportTwo.pdf](https://www.resource-recycling.com/images/BANReportTwo.pdf), accessed 30.05.2017).

772 Hu, S., Zhang, W., Li, J., Lin, K., Ji, R., 2016. Antioxidant and gene expression responses of *Eisenia fetida*  
 773 following repeated exposure to BDE209 and Pb in a soil-earthworm system. *Sci. Total Environ.* 556, 163-  
 774 168.

775 Huang, Y., Ni, W., Chen, Y., Wang, X., Zhang, J., Wu, K., 2015. Levels and risk factors of antimony  
 776 contamination in human hair from an electronic waste recycling area, Guiyu, China. *Environ. Sci. Pollut.*  
 777 *Res.* 22(9), 7112-7119.

778 Huisman, J., Botezatu, I., Herreras, L., Liddane, M., Hintsa, J., Luda di Cortemiglia, et al., 2015.  
 779 Countering WEEE Illegal Trade (CWIT) Summary Report, Market Assessment, Legal Analysis, Crime  
 780 and Recommendations Roadmap, August 30, 2015. Lyon, France.  
 781 (<https://www.interpol.int/content/download/32151/417087/version/1/file/CWIT%203%20small.pdf>,  
 782 accessed 30.05.2017).

783 IARC (International Agency for Research on Cancer), IARC Monographs on the Evaluation of  
 784 Carcinogenic Risks to Humans (<http://monographs.iarc.fr/index.php>, accessed 30.05.2017)

785 Iannicelli-Zubiani, E. M., Giani, M. I., Recanati, F., Dotelli, G., Puricelli, S., Cristiani, C., 2017.  
 786 Environmental impacts of a hydrometallurgical process for electronic waste treatment: A life cycle  
 787 assessment case study. *J. Clean. Prod.* 140, Part 3, 1204-1216.

788 Interpol, 2009. Electronic Waste and Organized Crime - Assessing the link - Phase II Report for the  
 789 Interpol Pollution Crime Working Group: Interpol Pollution Crime Working Group  
 790 (<https://www.interpol.int/content/download/5367/45070/version/3/file/Wastereport.pdf>,  
 791 accessed 30.05.2017).

792 Iqbal, M., Breivik, K., Syed, J. H., Malik, R. N., Li, J., Zhang, G., Jones, K. C., 2015. Emerging issue of  
 793 e-waste in Pakistan: A review of status, research needs and data gaps. *Environ. Pollut.* 207, 308-318.

794 Julander, A., Lundgren, L., Skare, L., Grandér, M., Palm, B., Vahter, M., Lidén, C., 2014. Formal  
 795 recycling of e-waste leads to increased exposure to toxic metals: An occupational exposure study from  
 796 Sweden. *Environ. Int.* 73, 243-251.

797 Kiddee, P., Naidu, R., Wong, M. H., 2013. Electronic waste management approaches: an overview. *Waste*  
 798 *Manag.* 33(5), 1237-1250.

799 Klaassen C., 2013. *Casarett & Doull's Toxicology: The Basic Science of Poisons*, Eighth Edition:  
 800 McGraw-Hill Education.

801 Klees, M., Hombrecher, K., Gladtko, D., 2017. Polychlorinated biphenyls in the surrounding of an e-waste  
 802 recycling facility in North-Rhine Westphalia: Levels in plants and dusts, spatial distribution, homologue  
 803 pattern and source identification using the combination of plants and wind direction data. *Sci. Total*  
 804 *Environ.* 603–604, 606-615.

805 Krieger, N., 2010. Workers are people too: societal aspects of occupational health disparities—an  
 806 ecosocial perspective. *Am. J. Ind. Med.* 53(2), 104–115.

807 Kumar, A., Holuszko, M., Espinosa, D. C. R., 2017. E-waste: An overview on generation, collection,  
 808 legislation and recycling practices. *Res. Conserv. Recy.* 122, 32-42.

809 Kumari, A., Jha, M. K., Lee, J.-C., Singh, R. P., 2016. Clean process for recovery of metals and recycling  
 810 of acid from the leach liquor of PCBs. *J. Clean. Prod.* 112, Part 5, 4826-4834.

811 Labunska, I., Abdallah, M. A.-E., Eulaers, I., Covaci, A., Tao, F., Wang, M., et al., 2015. Human dietary  
 812 intake of organohalogen contaminants at e-waste recycling sites in Eastern China. *Environ. Int.* 74, 209-  
 813 220.

814 Lebel, S., 2016. Fast machines, slow violence: ICTs, planned obsolescence, and e-waste. *Globalizations*.  
 815 13(3), 300-309.

816 Lecler, M. T., Zimmermann, F., Silvente, E., Clerc, F., Chollot, A., Grosjean, J., 2015. Exposure to  
817 hazardous substances in Cathode Ray Tube (CRT) recycling sites in France. *Waste Manag.* 39, 226-235.

818 Lepawsky, J., 2015. The changing geography of global trade in electronic discards: time to rethink the e-  
819 waste problem. *Geog. J.* 181(2), 147-159.

820 Leong, F. T. L., Eggerth, D.E., Chang, D., Flynn, M.A., Ford, J.K., Martinez, R.O., 2017. Introduction.  
821 In: Leong, F. T.L., Eggerth, D.E., Chang, D., Flynn, M.A., Ford, J. K., Martinez, R.O. (Eds.).  
822 Occupational Health Disparities among Racial and Ethnic Minorities: Formulating Research Needs and  
823 Directions. American Psychological Association. Washington, D.C.

824 Li, J., Zeng, X., Chen, M., Ogunseitan, O. A., Stevels, A., 2015. "Control-alt-delete": rebooting solutions  
825 for the E-waste problem. *Environ. Sci. Technol.* 49(12), 7095-7108.

826 Li, J., He, X., Xianlai Zeng., 2016. Designing and examining e-waste recycling process: methodology and  
827 case studies. *Environ. Tech.* DOI:10.1080/09593330.2016.1207711

828 Li, X., Tian, Y., Zhang, Y., Ben, Y., Lv, Q., 2017. Accumulation of polybrominated diphenyl ethers in  
829 breast milk of women from an e-waste recycling center in China. *J. Environ. Sci.* 52, 305-313.

830 Liang, L., Sharp, A., 2016. Determination of the knowledge of e-waste disposal impacts on the  
831 environment among different gender and age groups in China, Laos, and Thailand. *Waste Manag. Res.*  
832 34(4), 388-395.

833 Lin, Y., Xu, X., Dai, Y., Zhang, Y., Li, W., Huo, X., 2016. Considerable decrease of antibody titers against  
834 measles, mumps, and rubella in preschool children from an e-waste recycling area. *Sci. Tot. Environ.* 573,  
835 760-766.

836 Lin, X., Xu, X., Zeng, X., Xu, L., Zeng, Z., Huo, X., 2017. Decreased vaccine antibody titers following  
837 exposure to multiple metals and metalloids in e-waste-exposed preschool children. *Environ. Pollut.* 220.  
838 Pt A, 354-363.

839 Lines, K., Graside, B., Sinha, S., Fedorenko, I., 2016. Clean and inclusive? Recycling e-waste in China  
 840 and India: International Institute for Environment and Development  
 841 (<http://pubs.iied.org/pdfs/16611IIED.pdf>, accessed 30.05.2017).

842 Lu, S. Y., Li, Y. X., Zhang, T., Cai, D., Ruan, J. J., Huang, M. Z., et al., 2017. Effect of E-waste recycling  
 843 on urinary metabolites of organophosphate flame retardants and plasticizers and their association with  
 844 oxidative stress. *Environ. Sci. Tech.* 51 (4), 2427-2437.

845 Lundgren, K., 2012. The Global Impact of E-waste: Addressing the Challenge. International Labour  
 846 Organization, Geneva, Switzerland ([http://ilo.org/sector/Resources/publications/WCMS\\_196105/lang--](http://ilo.org/sector/Resources/publications/WCMS_196105/lang--en/index.htm)  
 847 [en/index.htm](http://ilo.org/sector/Resources/publications/WCMS_196105/lang--en/index.htm), accessed 30.05.2017)

848 Ma, J., Kannan, K., Cheng, J., Hori, Y., Wu, Q., Wang, W., 2008. Concentrations, profiles, and estimated  
 849 human exposures for polychlorinated dibenzo-p-dioxins and dibenzofurans from electronic waste  
 850 recycling facilities and a chemical industrial complex in Eastern China, *Environ. Sci. Technol.*, 42, 8252–  
 851 8259

852 Ma, J., Addink, R., Yun, S., Cheng, J., Wang, W., Kannan, K., 2009. Polybrominated dibenzo-p-  
 853 dioxins/dibenzofurans and polybrominated diphenyl ethers in soil, vegetation, workshop-floor dust, and  
 854 electronic shredder residue from an electronic waste recycling facility and in soils from a chemical  
 855 industrial complex in Eastern China. *Environ. Sci. Technol.* 43, 7350–7356.

856 Ma, Z. Q., Jung, Y. H., Chang, T. H., Seo, J. H., Zhang, H., Cai, Z., Gong, S., 2016. Green microwave  
 857 electronics for the coming era of flexible electronics. *Proceedings from Microwave Symposium (IMS),*  
 858 *IEEE MTT-S International*, San Francisco, CA, USA [22-27 May].

859 Mackintosh, S. A., Wallace, J. S., Gross, M. S., Navarro, D. D., Pérez-Fuentetaja, A., Alaei, M., et al.,  
 860 2015. Review on the occurrence and profiles of polybrominated diphenyl ethers in the Philippines.  
 861 *Environ. Int.* 85, 314-326.

862 Magalini, F., 2016. Global challenges for e-waste management: the societal implications. *Rev. Environ.*  
 863 *Health.* 31(1), 137-140.



864 Massari, M., Monzini, P., 2006. Dirty businesses in Italy: a case-study of illegal trafficking in hazardous  
865 waste. *Global Crime*. 6(3-4), 285-304.

866 Matsukami, H., Suzuki, G., Tue, N. M., Tuyen, L. H., Viet, P. H., Takahashi, S., et al., 2016. Analysis of  
867 monomeric and oligomeric organophosphorus flame retardants in fish muscle tissues using liquid  
868 chromatography–electrospray ionization tandem mass spectrometry: Application to Nile tilapia  
869 (*Oreochromis niloticus*) from an e-waste processing area in northern Vietnam. *Emerg. Contam.* 2(2), 89-  
870 97.

871 Matsukami, H., Suzuki, G., Someya, M., Uchida, N., Tue, N. M., Tuyen, L. H., et al., 2017. Concentrations  
872 of polybrominated diphenyl ethers and alternative flame retardants in surface soils and river sediments  
873 from an electronic waste-processing area in northern Vietnam, 2012–2014. *Chemosphere*, 167, 291-299.

874 McCann, D., Wittmann, A., 2015. E-waste Prevention, Take-back System Design and Policy Approaches.  
875 United Nations University/Step Initiative ([http://www.step-initiative.org/news/step-green-paper-](http://www.step-initiative.org/news/step-green-paper-explores-policy-solutions-to-e-waste-problem.html)  
876 [explores-policy-solutions-to-e-waste-problem.html](http://www.step-initiative.org/news/step-green-paper-explores-policy-solutions-to-e-waste-problem.html), accessed 30.05.2017).

877 Meyer, D. E., Katz, J. P., 2016. Analyzing the environmental impacts of laptop enclosures using screening-  
878 level life cycle assessment to support sustainable consumer electronics. *J. Clean. Prod.* 112, 369-383.

879 Milovantseva, N., Fitzpatrick, C., 2016. Effect of Waste Legislation on Transboundary Movements of  
880 EEE Destined for Reuse: Impact of E-waste Regulations on Reuse Organizations and Possible Future  
881 Direction. United Nations University/Step Initiative ([http://www.step-initiative.org/files/step-](http://www.step-initiative.org/files/step-2014/Publications/Green%20and%20White%20Papers/Step_GP_TBM_draft_fin_16_01_13.pdf)  
882 [2014/Publications/Green%20and%20White%20Papers/Step\\_GP\\_TBM\\_draft\\_fin\\_16\\_01\\_13.pdf](http://www.step-initiative.org/files/step-2014/Publications/Green%20and%20White%20Papers/Step_GP_TBM_draft_fin_16_01_13.pdf),  
883 accessed 3.06.2017)

884 MJC & Associates, 2004. Screening Level Human Health and Ecological Risk Assessment for Generic  
885 E-waste Processing Facility: ICT Innovation and Computers for Schools Program, Industry Canada and  
886 National Office of Pollution Prevention, Environment Canada. September 2004. MJC03-126.

887 Mohanty, S., Vermeersch, E., Hintsa, J., Di Cortemiglia, V. L., Liddane, M., 2015. Weaknesses in  
888 European e-Waste Management. Proceedings from the Hamburg International Conference of Logistics  
889 (HICL), Hamburg, Germany (<https://hicl.org/publications/2015/22/535.pdf>, accessed 10.06.2017)

890 Morris, A., Metternicht, G., 2016. Assessing effectiveness of WEEE management policy in Australia. J.  
891 Environ. Manag. 181, 218-230.

892 Movilla, A., N., Zwolinski, P., Dewulf, J., Mathieux, F., 2016. A method for manual disassembly analysis  
893 to support the ecodesign of electronic displays. Res. Conserv. Recyc. 114, 42-58.

894 Namias J., 2013. The future of electronic waste recycling in the United States: Obstacles and Domestic  
895 Solutions (MS thesis). Dept. Earth and Environmental Engineering, Columbia University, New York.

896 Nartey, K. V., 2016. Environmental and health impacts of informal e-waste recycling in Agbogbloshie,  
897 Accra, Ghana: Recommendations for Sustainable Management (PhD Thesis), Rheinischen Friedrich-  
898 Wilhelms-Universität Bonn, Germany.

899 NIOSH/ASSE (National Institute for Occupational Safety and Health/American Society for Safety  
900 Engineers), 2015. Overlapping vulnerabilities: the occupational safety and health of young workers in  
901 small construction firms. By Flynn MA, Cunningham TR, Guerin RJ, Keller B, Chapman LJ, Hudson D,  
902 Salgado C. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control  
903 and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No.  
904 2015-178.

905 O'Connor, M. P., Zinnerman, J. B., Anastas, P. T., Plata, D. L., 2016. A strategy for material supply chain  
906 sustainability: enabling a circular economy in the electronics industry through green engineering. ACS  
907 Sustainable Chem. Eng. 4(11), 5879-5888.

908 Page, E., Ceballos, D., Oza, A., Gong, W., Mueller, C., 2015. Metal Exposures in an Electronic Scrap  
909 Recycling Facility. U.S. Department of Health and Human Services, Centers for Disease Control and  
910 Prevention, National Institute for Occupational Safety and Health, Report No. 2013-0067-3228  
911 (<https://www.cdc.gov/niosh/hhe/reports/pdfs/2013-0067-3228.pdf>, accessed 06.08.2017).

912 Pascale, A, Sosa, A., Bares, C., Battocletti, A., Moll, M. J., Pose, D., et al., 2016. E-waste informal  
913 recycling: an emerging source of lead exposure in South America. Ann. Glob. Health. 82(1), 197-201.

914 Peluola, A., 2016. Investigation of the implementation and effectiveness of electronic waste management  
915 in Nigeria. *Model. Earth Syst. Environ.* 2:100.

916 Perkins, D. N., Brune Drisse, M.-N., Nxele, T., Sly, P. D., 2014. E-waste: a global hazard. *Ann. Glob.*  
917 *Health.* 80(4), 286-295.

918 Petridis, N. E., Stiakakis, E., Petridis, K., Dey, P., 2016. Estimation of computer waste quantities using  
919 forecasting techniques. *J. Clean. Prod.* 112, 3072-3085.

920 Pickren, G., 2015. Making connections between global production networks for used goods and the realm  
921 of production: a case study on e-waste governance. *Glob. Netw.* 14(4), 403-423.

922 Premalatha, M., Thabassum-Abbasi, Tasneem Abbasi, Abbasi, S.A., 2014. The generation, impact, and  
923 management of e-waste: State of the Art. *Crit. Rev. Env. Sci. Technol.* 44(14), 1577-1678.

924 Robinson, B.H., 2009. E-waste: An Assessment of global production and environmental impacts. *Sci.*  
925 *Total Environ.* 408 (2), 183-191.

926 Rucevska I., N. C., Isarin N., Yang W., Liu N., Yu K., Sandnæs S., et al., 2015. Waste Crime Waste Risks  
927 - Gaps in the Meeting the Global Waste Challenge, A UNEP Rapid Response Assessment. United Nations  
928 Environment Programme and GRID-Arendal, Nairobi and Arendal  
929 ([http://www.unep.org/ourplanet/september-2015/unep-publications/waste-crime-waste-risks-gaps-](http://www.unep.org/ourplanet/september-2015/unep-publications/waste-crime-waste-risks-gaps-meeting-global-waste-challenge-rapid)  
930 [meeting-global-waste-challenge-rapid](http://www.unep.org/ourplanet/september-2015/unep-publications/waste-crime-waste-risks-gaps-meeting-global-waste-challenge-rapid), accessed 01.06.2017).

931 Ryneal, M., 2016. Turning waste into gold: Accumulation by disposal and the political economy of e-  
932 waste urban mining (PhD Thesis), Northern Arizona University, Arizona, USA. ProQuest Number:  
933 10125349.

934 Sabbaghi, M., Cade, W., Behdad, S., Bisantz, A. M., 2017. The current status of the consumer electronics  
935 repair industry in the U.S.: a survey-based study. *Res. Conserv. Recy.* 116, 137-151.

936 Sahle-Demessie, E., Richardson, T., Changseok Han, C., Dietrich, J., Wang, J., 2016. Characterizing of  
937 Emissions from Open Burning of Electronic Waste using TG-GC-MS System (<https://resource->

938 recycling.com/e-scrap/wp-content/uploads/sites/2/2017/02/Emissions-from-Open-Burning-of-  
 939 Electronic-Waste-Application-Note-013098\_01-2.pdf, accessed 02.06.2017).

940 Scruggs, C. E., Nimpuno, N., Moore, R. B. B., 2016. Improving information flow on chemicals in  
 941 electronic products and E-waste to minimize negative consequences for health and the environment.  
 942 Resources, Conservation and Recycling, 113, 149-164.

943 Schluep, M., Hagelueken, C., Kuehr, R., Magalini, F., Maurer, C, Meskers, E, et al., 2009. Recycling from  
 944 e-Waste to Resources. Sustainable Innovation and Technology Transfer Industrial Sector Studies.  
 945 UNEP/Step ([http://www.unep.org/pdf/Recycling\\_From\\_e-waste\\_to\\_resources.pdf](http://www.unep.org/pdf/Recycling_From_e-waste_to_resources.pdf), accessed  
 946 02.06.2017).

947 Seeberger, J., Grandhi, R, Kim, SS, Mase, WA, Reponen, T, Ho, S, Chen, A., 2016. E-waste management  
 948 in the United States and public health implications. J. Environ. Health. 79(3).

949 Shen, C., Tanga, X., Cheemaa, S. A., Zhanga, C., Khana, M. I., Lianga, F., et al., 2009. Enhanced  
 950 phytoremediation potential of polychlorinated biphenyl contaminated soil from e-waste recycling area in  
 951 the presence of randomly methylated- $\beta$ -cyclodextrins. J. Hazard. Mat. 172 (2-3), 1671-1676.

952 Shen, H., Guan, R., Ding, G., Chen, Q., Lou, X., Chen, Z., et al., 2017. Polychlorinated dibenzo-p-  
 953 dioxins/furans (PCDD/Fs) and polychlorinated biphenyls (PCBs) in Zhejiang foods (2006–2015): Market  
 954 basket and polluted areas. Sci. Tot. Environ. 574, 120-127.

955 Shi, J. C., Zheng, G. J. S., Wong, M. H., Liang, H., Li, Y. L., Wu, Y. L., et al., 2016. Health risks of  
 956 polycyclic aromatic hydrocarbons via fish consumption in Haimen bay (China), downstream of an e-waste  
 957 recycling site (Guiyu). Environ. Res. 147, 233-240.

958 Singh, N., Li, J., Zeng, X., 2016. Global responses for recycling waste CRTs in e-waste. Waste Manage.  
 959 57, 187-197.

960 Someya, M., Suzuki, G., Ionas, A. C., Tue, N. M., Xu, F., Matsukami, H., et al., 2016. Occurrence of  
 961 emerging flame retardants from e-waste recycling activities in the northern part of Vietnam. Emerg.  
 962 Contam. 2(2), 58-65.

963 Son, K.B., Lee, D. S., Lim, S.R., 2015. Effect of technology convergence for tablet PC on potential  
 964 environmental impacts from heavy metals. *Int. J. Sust. Dev. World* 23(2), 154-162.

965 Song, Q., Li, J., 2015. A review on human health consequences of metals exposure to e-waste in China.  
 966 *Environ. Pollut.* 196, 450-461.

967 Song, Q., Zeng, X., Li, J., Duan, H., Yuan, W., 2015. Environmental risk assessment of CRT and PCB  
 968 workshops in a mobile e-waste recycling plant. *Environ. Sci. Pollut. Res.* 22(16), 12366-12373.

969 Srighboh, R. K., Basu, N., Stephens, J., Asampong, E., Perkins, M., Neitzel, R. L., Fobil, J., 2016. Multiple  
 970 elemental exposures amongst workers at the Agbogbloshie electronic waste (e-waste) site in Ghana.  
 971 *Chemosphere*, 164, 68-74.

972 Step, 2014. One Global Definition of E-waste. Solving the E-Waste Problem (Step) White Paper Series.  
 973 United Nations University/Step Initiative 2014. ([http://www.step-](http://www.step-initiative.org/files/step/_documents/StEP_WP_One%20Global%20Definition%20of%20E-waste_20140603_amended.pdf)  
 974 [initiative.org/files/step/\\_documents/StEP\\_WP\\_One%20Global%20Definition%20of%20E-](http://www.step-initiative.org/files/step/_documents/StEP_WP_One%20Global%20Definition%20of%20E-waste_20140603_amended.pdf)  
 975 [waste\\_20140603\\_amended.pdf](http://www.step-initiative.org/files/step/_documents/StEP_WP_One%20Global%20Definition%20of%20E-waste_20140603_amended.pdf), accessed 06.08.2017).

976 Step, 2016. Guiding Principles to Develop E-waste Management Systems and Legislation. Solving the E-  
 977 Waste Problem (Step) White Paper Series. United Nations University/Step Initiative 2016.  
 978 ([http://www.step-initiative.org/files/step-](http://www.step-initiative.org/files/step-2014/Publications/Green%20and%20White%20Papers/Step_WP_WEEE%20systems%20and%20legislation_final.pdf)  
 979 [2014/Publications/Green%20and%20White%20Papers/Step\\_WP\\_WEEE%20systems%20and%20legisl](http://www.step-initiative.org/files/step-2014/Publications/Green%20and%20White%20Papers/Step_WP_WEEE%20systems%20and%20legislation_final.pdf)  
 980 [ation\\_final.pdf](http://www.step-initiative.org/files/step-2014/Publications/Green%20and%20White%20Papers/Step_WP_WEEE%20systems%20and%20legislation_final.pdf), accessed 30.05.2017).

981 Sthiannopkao, S., Wong, M. H., 2013. Handling e-waste in developed and developing countries:  
 982 Initiatives, practices, and consequences. *Sci. Tot. Environ.* 463–464, 1147-1153.

983 Sugimura, Y., Murakami, S., 2016. Problems in Japan's governance system related to end-of-life electrical  
 984 and electronic equipment trade. *Resour. Conserv. Recy.* 112, 93-106.

985 Suzuki, G., Someya, M., Matsukami, H., Tue, N. M., Uchida, N., Tuyen, L. H., et al., 2016.  
 986 Comprehensive evaluation of dioxins and dioxin-like compounds in surface soils and river sediments from

987 e-waste-processing sites in a village in northern Vietnam: Heading towards the environmentally sound  
 988 management of e-waste. *Emerg. Contam.* 2(2), 98-108.

989 Szalatkiewicz, J., 2014. Metals content in printed circuit board waste. *Pol. J. Environ. Stud.*, 23(6), 2365-  
 990 2369.

991 Tansel, B., 2017. From electronic consumer products to e-wastes: Global outlook, waste quantities,  
 992 recycling challenges. *Environ. Int.* 98, 35-45.

993 Tao, F., Matsukami, H., Suzuki, G., Nguyen Minh, T., Pham Hung, V., Takigami, H., Harrad, S., 2016.  
 994 Emerging halogenated flame retardants and hexabromocyclododecanes in food samples from an e-waste  
 995 processing area in Vietnam. *Environ. Sci. Proc. Imp.* 18, 361-370.

996 Terazono, A., Oguchi, M., Yoshida, A., Medina, R. P., Ballesteros, F. C. Jr., 2017. Material recovery and  
 997 environmental impact by informal e-waste recycling site in the Philippines. In Matsumoto et al. (Eds.),  
 998 Sustainability through Innovation in Product Life Cycle Design, Vol. Part II, Singapore: Springer  
 999 Singapore, pp. 197-213.

1000 Toxnet (n.d.). Toxicology Data Network. (<https://toxnet.nlm.nih.gov/cgi-bin/sis/search2>, accessed  
 1001 30.05.2017).

1002 Tran, H. P., Wang, F., Dewel, J., Huynh, T. H., Schaubroeck, T., 2016. Estimation of the unregistered  
 1003 inflow of electrical and electronic equipment to a domestic market: A case study on televisions in Vietnam.  
 1004 *Environ. Sci. Technol.* 50 2424–2433.

1005 Tue, N. M., Goto, A., Takahashi, S., Itai, T., Asante, K. A., Nomiya, K., et al., 2016a. Soil  
 1006 contamination by halogenated polycyclic aromatic hydrocarbons from open burning of e-waste in  
 1007 Agbogbloshie (Accra, Ghana). *J. Mat. Cycles Waste Manag.* DOI: 10.1007/s10163-016-0568-y

1008 Tue, N. M., Goto, A., Takahashi, S., Itai, T., Asante, K. A., Kunisue, T., Tanabe, S., 2016b. Release of  
 1009 chlorinated, brominated and mixed halogenated dioxin-related compounds to soils from open burning of  
 1010 e-waste in Agbogbloshie (Accra, Ghana). *J. Hazard. Mater.* 302, 151-157.

1011 UNEP (United Nations Environment Programme), 2010. Introductory Note to the Basel Convention by  
 1012 Dr. Katharina Kummer Peiry, Executive Secretary of the Basel Convention, UNEP  
 1013 (<http://legal.un.org/avl/ha/bcctmhwd/bcctmhwd.html>, accessed 15 June 2017).

1014 US DOJ (United States Department of Justice), 2010. A Review of Federal Prison Industries' Electronic-  
 1015 Waste Recycling Program. Office of the Inspector General, Oversight and Review Division; October  
 1016 2010.

1017 US EPA (United States Environmental Protection Agency), 2015. Flame Retardants in Printed Circuit  
 1018 Boards. Chapter 3. Final Report. EPA Publication 744-R-15-001.  
 1019 ([https://www.epa.gov/sites/production/files/2015-08/documents/pcb\\_final\\_report.pdf](https://www.epa.gov/sites/production/files/2015-08/documents/pcb_final_report.pdf), accessed  
 1020 15.06.2017).

1021 US EPA, 2016a. EPD Will Not Tolerate Illegal Import Of Hazardous E-Waste, Press Release, June 17,  
 1022 2016. (<http://www.info.gov.hk/gia/general/201606/17/P201606171114.htm>, accessed 15.06.2017).

1023 US EPA, 2016b. Implementation Study Of The Electronics Recycling Standards: R2 and e-Stewards®:  
 1024 United States Environmental Protection Agency, Office of Resource Conservation and Recovery and Abt  
 1025 Associates, Inc.

1026 Verdecchia, R., Ricchiuti, F., Hankel, A., Lago, P., Procaccianti, G., 2017. Green ICT research and  
 1027 challenges. In V. Wohlgemuth, F. Fuchs-Kittowski & J. Wittmann (Eds.), *Advances and New Trends in*  
 1028 *Environmental Informatics: Stability, Continuity, Innovation*, pp. 37-48. Cham: Springer International  
 1029 Publishing.

1030 Wang, Yan, Luo, C., Wang, S., Cheng, Z., Li, J., Gan, Z., 2016. The abandoned e-waste recycling site  
 1031 continued to act as a significant source of polychlorinated biphenyls: An in situ assessment using fugacity  
 1032 samplers. *Environ. Sci. Technol.*, 50, 8623–8630.

1033 Wang, Yan, Wu, X., Hou, M., Zhao, H., Chen, R., Luo, C., Zhang, G., 2017. Factors influencing the  
 1034 atmospheric concentrations of PCBs at an abandoned e-waste recycling site in South China. *Sci. Tot.*  
 1035 *Environ.* 578, 34-39.



- 1036 Wang, Yalin, Hu, J., Lin, W., Wang, N., Li, C., Luo, et al. 2016. Health risk assessment of migrant  
1037 workers' exposure to polychlorinated biphenyls in air and dust in an e-waste recycling area in China:  
1038 Indication for a new wealth gap in environmental rights. *Environ. Int.* 87, 33-41.
- 1039 Wilson D. C., Rodic., L., Modak, P., Soos, R., Carpintero Rogero, A., Velis, C., et al. 2015. Global Waste  
1040 Management Outlook. UNEP. ([http://www.unep.org/ourplanet/september-2015/unep-](http://www.unep.org/ourplanet/september-2015/unep-publications/global-waste-management-outlook)  
1041 [publications/global-waste-management-outlook](http://www.unep.org/ourplanet/september-2015/unep-publications/global-waste-management-outlook), accesses 30.05.2017).
- 1042 Wilson, G. T., Smalley, G., Suckling, J. R., Lilley, D., Lee, J., Mawle, R. 2017. The hibernating mobile  
1043 phone: Dead storage as a barrier to efficient electronic waste recovery. *Waste Manag.* 60, 521-533.
- 1044 WDE (Washington State Department of Ecology), 2016. Electronics Recycler Fined \$444,000 for Illegally  
1045 Disposing of Dangerous Waste Overseas, 1 September 2016.  
1046 (<http://www.ecy.wa.gov/news/2016/114.html>, accessed 30.05.2017).
- 1047 Wittsiepe, J., Fobil, J. N., Till, H., Burchard, G.-D., Wilhelm, M., Feldt, T., 2015. Levels of  
1048 polychlorinated dibenzo-p-dioxins, dibenzofurans (PCDD/Fs) and biphenyls (PCBs) in blood of informal  
1049 e-waste recycling workers from Agbogbloshie, Ghana, and controls. *Environ. Int.* 79, 65-73.
- 1050 Woo, S. H., Lee, D. S., Lim, S. R., 2016. Potential resource and toxicity impacts from metals in waste  
1051 electronic devices. *Integr. Environ. Asses.* 12(2), 364-370.
- 1052 Wu, Chen-Chou, Bao, L.-J., Tao, S., Zeng, E. Y., 2016. Dermal uptake from airborne organics as an  
1053 important route of human exposure to e-waste combustion fumes. *Environ. Sci. Technol.* 50(13), 6599-  
1054 6605.
- 1055 Wu, Q., Leung, J. Y. S., Geng, X., Chen, S., Huang, X., Li, H., et al., 2015. Heavy metal contamination  
1056 of soil and water in the vicinity of an abandoned e-waste recycling site: Implications for dissemination of  
1057 heavy metals. *Sci. Tot. Environ.* 506-507, 217-225.
- 1058 Wu, Ping-Fan, Yu, L. L., Li, L., Zhang, Y., Li, X. H., 2016. Maternal transfer of dechloranes and their  
1059 distribution among tissues in contaminated ducks. *Chemosphere*, 150, 514-519.



- 1060 Xiao, X., Hu, J., Peng, P. a., Chen, D., Bi, X., 2016. Characterization of polybrominated dibenzo-p-dioxins  
1061 and dibenzo-furans (PBDDs/Fs) in environmental matrices from an intensive electronic waste recycling  
1062 site, South China. *Environ. Pollut.* 212, 464-471.
- 1063 Xu, Long, Huo, X., Zhang, Y., Li, W., Zhang, J., Xu, X., 2015. Polybrominated diphenyl ethers in human  
1064 placenta associated with neonatal physiological development at a typical e-waste recycling area in China.  
1065 *Environ. Pollut.* 196, 414-422.
- 1066 Xu, Long, Ge, J., Huo, X., Zhang, Y., Lau, A. T. Y., Xu, X., 2016. Differential proteomic expression of  
1067 human placenta and fetal development following e-waste lead and cadmium exposure in utero. *Sci. Tot.*  
1068 *Environ.* 550, 1163-1170.
- 1069 Xu, Pengjun, Tao, B., Ye, Z., Zhao, H., Ren, Y., Zhang, T., et al., 2016. Polycyclic aromatic hydrocarbon  
1070 concentrations, compositions, sources, and associated carcinogenic risks to humans in farmland soils and  
1071 riverine sediments from Guiyu, China. *J. Environ. Sci.* 48, 102-111.
- 1072 Xu, Xinjin, Zeng, X., Boezen, H. M., Huo, X., 2015. E-waste environmental contamination and harm to  
1073 public health in China. *Front. Med-PRC.* 9(2), 220-228.
- 1074 Yadav, V., Cheraghi, S. A., Namboodiri, V., 2015. The OS elixir: On the impact of operating system  
1075 upgrades on mobile device lifespan. *Proceeding from Green Computing Conference and Sustainable*  
1076 *Computing Conference (IGSC), 2015 Sixth International, Las Vegas, NV, USA, [14-16 December].*
- 1077 You, Z. W., Hou, W. J., Wang, X. C., Ma, J. S., Liu, J., 2016. Design strategies for developing sustainable  
1078 consumer electronics. *J. Residual. Sci. Tech.* 13, S219-S226.
- 1079 Yuan, B., Fu, J., Wang, Y., Jiang, G., 2017. Short-chain chlorinated paraffins in soil, paddy seeds (*Oryza*  
1080 *sativa*) and snails (*Ampullariidae*) in an e-waste dismantling area in China: Homologue group pattern,  
1081 spatial distribution and risk assessment. *Environ. Pollut.*
- 1082 Yu-Gong, Tian, X.-m., Wu, Y.-f., Zhe-Tan, Lei-Lv, 2016. Recent development of recycling lead from  
1083 scrap CRTs: a technological review. *Waste Manag.* 57, 176-186.

1084 Zeng, Xiang, Xu, X., Boezen, H. M., Huo, X., 2016. Children with health impairments by heavy metals  
1085 in an e-waste recycling area. *Chemosphere*, 148, 408-415.

1086 Zeng, Xianlai, Gong, R., Chen, W.Q, Jinhui, I., 2016. Uncovering the recycling potential of “new” WEEE  
1087 in China. *Environ. Sci. Technol.*, 50, 1347–1358.

1088 Zeng, X., Yang, C., Chiang, J. F., Li, J., 2017. Innovating e-waste management: From macroscopic to  
1089 microscopic scales. *Sci. Total Environ.* 575, 1-5.

1090 Zhao, Y., Ma, J., Qiu, X., Lin, Y., Yang, Q., Zhu., T., 2013. Gridded field observations of polybrominated  
1091 diphenyl ethers and decabromodiphenyl ethane in the atmosphere of North China, *Environ. Sci. Technol.*  
1092 47(15), 8123- 8129.

1093 Zhang, B., Guan, D., 2016. Take responsibility for electronic waste disposal. *Nature*. 536 (7614), 23-25.

1094 Zhang, L., Xu, Z., 2016. A review of current progress of recycling technologies for metals from waste  
1095 electrical and electronic equipment. *J. Clean. Prod.* 127, 19-36.

1096 Zhang, Tao, Xue, J., Gao, C. Z., Qiu, R. L., Li, Y. X., Li, X., et al., 2016. Urinary concentrations of  
1097 bisphenols and their association with biomarkers of oxidative stress in people living near e-waste recycling  
1098 facilities in China. *Environ. Sci. Technol.* 50(7), 4045-4053.

1099 Zheng, Xiaobo, Xu, F., Chen, K., Zeng, Y., Luo, X., Chen, S., et al., 2015. Flame retardants and  
1100 organochlorines in indoor dust from several e-waste recycling sites in South China: Composition  
1101 variations and implications for human exposure. *Environ. Int.* 78, 1-7.

1102 Zheng, Xiaobo, Xu, F., Luo, X., Mai, B., Covaci, A., 2016. Phosphate flame retardants and novel  
1103 brominated flame retardants in home-produced eggs from an e-waste recycling region in China.  
1104 *Chemosphere*. 150, 545-550.

1105 Zheng, Jing, Yu, L. H., Chen, S. J., Hu, G. C., Chen, K. H., Yan, X., et al., 2016. Polychlorinated Biphenyls  
1106 (PCBs) in human hair and serum from e-waste recycling workers in Southern China: Concentrations,  
1107 chiral signatures, correlations, and source identification. *Environ. Sci. Technol.* 50(3), 1579-1586.

- 1108 Zheng, Xiangbin, Xu, X., Yekeen, T. A., Zhang, Y., Chen, A., Kim, S. S., et al., 2016. Ambient air heavy  
1109 metals in PM 2.5 and potential human health risk assessment in an informal electronic-waste recycling  
1110 site of China. *Aerosol Air. Qual. Res.* 16, 388-397.
- 1111 Zhu, C., Wang, P., Li, Y., Chen, Z., Li, H., Ssebugere, P., Zhang, Q., Jiang, Q., 2017. Trophic transfer of  
1112 hexabromocyclododecane in the terrestrial and aquatic food webs from an e-waste dismantling region in  
1113 East China. *Environ. Sci.: Processes Impacts*.

DRAFT

## APPENDIX

**Table 1**

### Recent reports on environmental and human burdens ascribed to e-waste contaminants

Nature of activities, sampling site	Environmental matrices sampled and dates	Targeted contaminants	Reported outcomes	Human health assessment	References
Informal small dismantling workshops and factories in the city of Guiyu <sup>1</sup> (China)	Topsoil and riverine sediment within Guiyu and adjacent towns February 2009	PAHs	High concentration of PAHs compared to rural soils elsewhere in China. Concentrations of PAHs higher in sediments compared to soils.	Relatively high carcinogenic risks associated with PAHs in riverine sediments through soil ingestion, dermal absorption and particulates inhalation.	Xu Pengjun et al., 2016
	Air in the vicinity of open burning and crude recycling March 2012 to April 2013	Heavy metals	Larger PM <sub>2.5</sub> concentrations than in other Asian cities, with higher heavy metal concentrations during winter and spring.	- Total potential cancer risk for both adults and children higher than US EPA recommendations. - Higher public health risk for children than adults, with regard to carcinogenic and non-carcinogenic elements in PM <sub>2.5</sub> .	Zheng Xiangbin et al., 2016
Informal e-recycling industrial park, in a collection of small villages around Taizhou (Zhejiang, China)	Soil, paddy seeds (hulls and unpolished rice), apple snails. Sampling area of 65-km radius around Taizhou. 2008 and 2010	Short-chain chlorinated paraffins (SCCPs) <sup>2</sup>	Higher concentrations in snails and rice grain hulls (part of the feed formulation of pigs and chickens on local farms) than in soil and unpolished rice.	Elevated risks via dietary exposure to SCCPs for people living in the e-recycling area.	Yuan et al., 2017
Informal e-recycling sites (Southern China)	Indoor dust September 2013 and March 2014	PCBs PBDEs DBDPE DPs Heavy metals	High concentrations of organic pollutants and heavy metals (except for Zn) in e-recycling areas.	- Excessive estimated daily intakes (EDIs) of PCBs, PBDEs, DBDPE, and DPs for adults in the e-waste recycling area (respectively 5.7–18.6 times and 4.1–85.5 times higher than those in rural and urban areas). - An EDI of Cd in the e-waste recycling 11.8 to 19.8 times higher than that in rural and urban areas. - An EDI of Pb 18 times higher than the Pb reference dose.	He et al., 2017
Informal e-recycling Zhejiang market (China)	620 foods collected between 2006-2015	PCDD/Fs PCBs	High concentrations of PCDD/Fs and PCBs in all foods sampled.	High human risk in the worst case scenario (only highly contaminated foods consumed). Estimated dietary intake for local residents approximatively 3.5-fold greater than the standard toxic equivalent (TEQ).	Shen et al., 2017
Informal e-recycling workshops and open	Soil in workshops and open burning areas	PAHs	Predominance of phenanthrene, fluoranthene and pyrene.	Elevated carcinogenic risk in case of prolonged exposure via multiple pathways	Daso et al., 2016

Nature of activities, sampling site	Environmental matrices sampled and dates	Targeted contaminants	Reported outcomes	Human health assessment	References
burning facilities Agbogbloshie (Ghana) (one of the world's largest e-recycling areas)	2015	PBDEs Heavy metals	Notable presence of PBDEs in surface soils; BDE-28 congener predominant. Moderate to extreme pollution by metals.	With moderate and maximum exposure scenarios via ingestion: significant non-carcinogenic health risk of local PBDE (BDE-47 and BDE-99 congeners) exposure for children and adverse health predicted for both adults and children with regard to Cu and Pb.	Akortia et al., 2017
Informal metal recovery or dismantling workshops, nearby highway in urban centers and suburban industrial roadsides in Chennai City (India)	Dust from workshops and nearby other sample points May 2014	PCBs	Concentration of PCBs in settled dust from informal metal recovery workshops tenfold higher than in urban centre.	- Major exposure routes for PCBs: dermal contact, ingestion (hand-to-mouth) and dust inhalation. - High cancer risk in e-recycling workshops for children via ingestion exposure. - Ingestion and dermal contact risk in e-recycling workshops exceeding Canadian threshold limit for a public health concern ( $1 \times 10^{-6}$ ).	Chakraborty et al., 2016
Informal e-recycling area Ogunpa Market, Ibadan (Nigeria)	Soil from different depths (within recycling and open burning sites) Unspecified period	Heavy metals and metalloids	Heavy contamination, with average concentrations of Pb and Cu exceeding soil regulatory limits. 65% of Cu and 88% of Pb susceptible to potential mobility.		Adesokan et al., 2016
Informal e-recycling site Meycauayan, Bulacan operating for >30 years (The Philippines)	Surface soil within e-recycling area January 2012	Heavy metals	High concentrations of Pb and Cu, especially near smelting area and stockyard.		Terazono et al., 2017
Informal e-recycling in workshops among residences Sue Yai Utit, Bangkok (Thailand)	Surface soil (within e-dismantling site and in adjacent areas) August 2014	Heavy metals	High contamination levels of Cu, Ni, Pb, and Zn, in both fine and coarse soil particles.		Damrongsiri et al., 2016
Backyard informal e-recycling areas, often within 20 m of living area Bui Dau village, Hung Yen province, (North Vietnam)	Fish (from ponds and canals close to workshops), sediment (drainage canals of water flow near families having workshops), house dust (surface of furniture and fan blades in the houses) 2013-2015	Flame retardants PBDEs	<i>Fish</i> : High concentrations of PBDEs compared to other fish collected in some farming facilities. <i>Sediments</i> : High concentrations of PBDEs compared to sediments collected from other industrial areas. <i>House dust</i> : High concentrations of PBDEs (BDE-99 and BDE-209) compared to house dust in Australia or Canada (but lower than indoor dust from e-waste recycling sites in China).	- Intakes of PBDEs, especially BDE-99 and BDE-209 congeners, higher via fish consumption than intakes via dust ingestion. - Higher range of body weight normalizing daily intake through dust ingestion than those reported in other countries.	Anh et al., 2016

Nature of activities, sampling site	Environmental matrices sampled and dates	Targeted contaminants	Reported outcomes	Human health assessment	References
The surroundings and the premises of formal e-recycling facilities in North-Rhine Westphalia, Germany	Curly kale and spruce needles from the residential area around the formal facility Street dust from or near the access roads to the facilities Dust from the premises 2014 and 2015	PCBs	<i>Curly kale and spruce needles:</i> High PCB concentrations directly north of the industrial premises. Evident concentration gradient from the industrial premises to the residential areas located southwest and northeast. <i>Street dust and dust:</i> High PCB contamination. Homologue patterns of highly PCB-contaminated dusts and street dusts similar to the homologue patterns of PCBs in curly kale and spruce needles.		Klees et al., 2017

<sup>1</sup> Informal activities have been suspended in Guiyu since December 2015 (Ceballos and Dong, 2016).

<sup>2</sup> The inclusion of SCCPs as persistent organic pollutants (POPs) in the Stockholm Convention is under review (Yuan et al., 2017).

**Table 2**

**Recent reports on the body burden of e-waste contaminants in e-recycling workers and surrounding populations**

Study population	Nature of activities, sampling site	Type of sample and date	Targeted contaminants	Reported outcomes	References
Workers	Informal e-recycling Longtang, Qingyuan City (China)	Hair and serum 2010	DPs	- Elevated serum and hair DPs levels. - Positive correlation between hair DPs levels and serum DPs levels in paired samples.	Chen et al., 2015
	Informal e-recycling (Southern China)	Hair and serum July 2011	PCBs	- Elevated serum and hair PCB levels. - Positive correlation between hair PCB levels and serum PCB levels in paired samples.	Zheng Jing et al., 2016
Surrounding population	Informal e-recycling Guiyu (China)	Hair from people living and/or working in the e-recycling area April-June 2012	Sb	- Higher Sb levels than at control site. - Higher Sb levels for people in e-recycling activities. - Main factors associated with high Sb levels: e-recycling work and long-term residence in Guiyu.	Huang et al., 2015
		Human placental tissues March and August 2012	PBDEs	Higher PBDE levels in placenta from women residing in Guiyu compared to control population.	Xu Long et al., 2015
			Cd, Pb	Placental Pb and Cd levels higher in the e-recycling area.	Xu Long et al., 2016
	Informal e-recycling Taizhou (China)	Children's blood 2012	Heavy metals and metalloids	High levels of Pb, Ni, Cu, Zn, and Se in exposed children compared to reference group living in a fishing town.	Lin et al., 2017
		Breast milk of women living within the e-recycling area From 2012 to 2013	PBDEs, hydroxylated polychlorinated biphenyls (OH-PBDEs), 2,2',4,4',5,6'-hexachlorobiphenyl (CB-153)	- High median levels of PBDEs in breast milk from residents living in e-recycling areas via dietary exposure. - Great concern about potential effect on development of neonates over long-term exposure.	Li et al., 2017
		Urine July-August 2014	Bisphenols A (BPA), F (BPF) and S (BPS)	Urinary levels of BPA and BPF significantly higher in residents of e-recycling area compared to reference rural residents.	Zhang et al., 2016
Workers	CRT televisions Formal recycling (France)	Personal air and ambient air in a CRT recycling unit 2007-2011	Heavy metals	- High levels of pollution exceeding some French Ministry of Labour occupational exposure limits - Personal Pb dust exposure levels higher than the ambient levels in sampled facilities.	Leclercq et al., 2015
Workers	Informal e-recycling Agbogbloshie (Ghana)	Blood October 2011	PCDD/Fs, PCBs	PCDD/Fs blood levels related to the e-waste burning activities.	Wittsiepe et al., 2015
		Blood 2014	Pb	Blood Pb levels over five times higher than average levels in the US.	Carlson, 2016
		Blood, urine April 2014	Heavy metals	- Blood Cd and Pb levels and urinary As level higher than in reference group.	Srigboh et al., 2016

Study population	Nature of activities, sampling site	Type of sample and date	Targeted contaminants	Reported outcomes	References
				<ul style="list-style-type: none"> <li>- Urinary As level above the US Agency for Toxic Substances and Disease Registry (ATSDR) limit.</li> <li>- Highest biomarker levels among workers involved in e-waste open burning.</li> </ul>	
Workers	Formal recycling (Sweden)	Blood, urine, plasma 2007-2009	Heavy metals	Higher blood, urine, and/or plasma levels of Cr, Co, Hg, In, Pb in recycling workers, compared to office workers.	Julander et al., 2014
Workers	Formal e-recycling Unspecified location (3 facilities in the United States)	Blood, urine, personal air, surface and skin wipes January, February, April, June, September 2013 and February 2014	Heavy metals	<ul style="list-style-type: none"> <li>- One exposure to Pb and two exposures to Cd above the respective Occupational Safety and Health Administration Permissible Exposure Limit (OSHA PEL)</li> <li>- High blood Pb levels for some dismantler employees</li> <li>- Presence of heavy metals on the skin and clothing of employees</li> </ul>	Ceballos et al., 2017
Surrounding population	Informal e-recycling (burning cables) in Montevideo (Uruguay)	Blood from children and adolescents involved in burning cables and gathering metals 2010-2014	Pb	<ul style="list-style-type: none"> <li>- Blood Pb levels higher than recommended level of 5 µg/dL (microgram per deciliter)</li> <li>- High blood lead levels among youngest children (often playing in the ground where cables are burned)</li> </ul>	Pascale et al., 2016
Workers	Informal e-recycling Bui Dau (Vietnam)	Serum 2010-2011	Organohalogen compounds: PCBs, OH-PCBs, PBDEs, methoxylated polybrominated diphenyl ether (MeO-PBDEs), OH-PBDEs, bromophenols (BPhs)	Higher serum residue levels of PCBs, OH-PCBs, PBDEs, and BPhs in e-recycling workers compared to blood donors from a rural site	Eguchi et al., 2015



DRAFT